Rhode Island College

Digital Commons @ RIC

Master's Theses, Dissertations, Graduate
Research and Major Papers Overview

Master's Theses, Dissertations, Graduate
Research and Major Papers

4-6-2020

A Comparison of the Efficacy of Mupirocin Versus Povidone-iodine in the Treatment of Nasal Colonization with MRSA/MSSA to Prevent Pos-operative Infections: An Integrative Review

Ashley O'Rourke

Follow this and additional works at: <https://digitalcommons.ric.edu/etd>

Recommended Citation

O'Rourke, Ashley, "A Comparison of the Efficacy of Mupirocin Versus Povidone-iodine in the Treatment of Nasal Colonization with MRSA/MSSA to Prevent Pos-operative Infections: An Integrative Review" (2020). *Master's Theses, Dissertations, Graduate Research and Major Papers Overview*. 362.
<https://digitalcommons.ric.edu/etd/362>

This Major Paper is brought to you for free and open access by the Master's Theses, Dissertations, Graduate Research and Major Papers at Digital Commons @ RIC. It has been accepted for inclusion in Master's Theses, Dissertations, Graduate Research and Major Papers Overview by an authorized administrator of Digital Commons @ RIC. For more information, please contact digitalcommons@ric.edu.

A COMPARISON OF THE EFFICACY OF MUPIROCIN VERSUS POVIDONE-
IODINE IN THE TREATMENT OF NASAL COLONIZATION WITH MRSA/MSSA
TO PREVENT POST-OPERATIVE INFECTIONS: AN INTEGRATIVE REVIEW

A Major Paper Presented

by

Ashley O'Rourke

Approved:

Committee Chairperson	_____	_____
		(Date)
Committee Members	_____	_____
		(Date)
	_____	_____
		(Date)
Director of Master's Program	_____	_____
		(Date)
Dean, School of Nursing	_____	_____
		(Date)

A COMPARISON OF THE EFFICACY OF MUPIROCIN VERSUS POVIDONE-
IODINE IN THE TREATMENT OF NASAL COLONIZATION WITH MRSA/MSSA
TO PREVENT POST-OPERATIVE INFECTIONS: AN INTEGRATIVE REVIEW

by

Ashley O'Rourke

A Major Paper Submitted in Partial Fulfillment

of the Requirements for the Degree of

Master of Science in Nursing

in

The School of Nursing

Rhode Island College

2020

Abstract

Surgical site infections are defined by the CDC as infections that occur after surgery in the part of the body that the surgery took place. Surgical site infections are now the costliest and most common hospital acquired infections. It is important for healthcare practitioners to prevent this costly and deadly surgical complication. The use of intranasal mupirocin has been widely used in healthcare institutions but due to concerns regarding cost, mupirocin resistance, compliance rates, and effectiveness, alternatives are being pursued. An important and promising alternative is intranasal povidone-iodine. The purpose of this integrative review was to compare the effectiveness of intranasal mupirocin versus intranasal povidone-iodine to treat nasal colonization with MSSA/MRSA to prevent post-operative surgical site infections. A search was completed including CINAHL, PubMed, and Google Scholar using keywords and inclusion and exclusion criteria, resulting in six studies being included. The selection of research articles was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram. Critical appraisal was conducted utilizing Polit and Beck's Guide to an Overall Critique of Qualitative Research, Quantitative Research Report, and Literature Reviews. Data collection for this review was performed using tables created specifically for this review. A cross-study analysis was conducted and summarized using descriptive data synthesis. Findings included that using intranasal povidone-iodine pre-operatively was as effective as intranasal mupirocin in the decolonization of MSSA/MRSA in the nares in the prevention of post-operative SSIs. More research is needed including further randomized control trials and larger studies, which the nurse practitioner can facilitate, research, educate, and promote to encourage policy change within their institution.

Acknowledgements

I would like to thank my husband, for without you I never would have made it this far. For all the times I wanted to give up, you encouraged me to fulfill my dream and keep going. You put in endless hours taking care of our young children to allow me time for my studies, never once complaining. You have been my rock and my biggest supporter for the past four years and for you, I am eternally grateful.

To Doctor Cynthia Padula, my first reader, I cannot thank you enough for all of your time, patience, and guidance in the completion of this major paper.

Table of Contents

Background/Statement of the Problem	1
Literature Review.....	3
Theoretical Framework	30
Method	32
Results	37
Summary and Conclusions	51
Recommendations and Implications for Advanced Nursing Practice	56
References	61
Appendices	69

A Comparison of the Efficacy of Mupirocin Versus Povidone-Iodine in the Treatment of Nasal Colonization with MRSA/MSSA to Prevent Post-Operative Infections:

An Integrative Review

Background/Statement of the Problem

Post-operative infections are an unfortunate complication after surgery that health care providers go to great lengths to prevent. There are many strategies used to prevent post-operative infections such as maintaining a sterile operating room and strict hand hygiene both during and after surgery. Despite the implementation of preventive strategies, post-operative infection remains a problem for a variety of reasons. One specific etiology relates to the fact that some people are colonized in their nostrils with bacteria which, left untreated, can invade the surgical site, causing a post-operative infection.

Two types of bacteria commonly colonized in the nostrils that have the potential to cause a post-operative infection are methicillin resistant staphylococcus aureus (MRSA) and methicillin sensitive staphylococcus aureus (MSSA). As much as 30% of the population are asymptomatically colonized with staphylococcus aureus (Sakr, Bregeon, Mege, Rolain, & Blin, 2018). Staphylococcus aureus can be found in various parts of the body including the skin, rectum, vagina, gastrointestinal tract, axilla, and the anterior nares. The anterior nares are the most common site of colonization (Sakr et al.). Nasal colonization with staphylococcus aureus can cause opportunistic and potentially life-threatening infections, such as surgical site infections (SSIs), that increase morbidity, mortality, and healthcare costs (Sakr et al.).

Patients colonized in the nares with MRSA have a 30% increased risk of developing a MRSA infection during their hospital stay (Sai, Laurent, Strale, Denis, & Byl, 2015). Therefore, it has become an important preventive strategy to test patients for a MRSA/MSSA infection before surgery, as decolonization is most effective in these populations that are only at risk for a short period of time (Septimus & Schweizer, 2016). Many people colonized in the nostrils with MRSA/MSSA are asymptomatic and thus unaware that they are carriers. Between 15-30% of healthy adults are nasally colonized with MSSA and 1-3% are colonized with MRSA; these patients have higher bacterial loads and are at an increased risk of developing an infection due to their colonization (Septimus & Schweizer).

Treatment with five days of mupirocin ointment to both nostrils before surgery to eradicate the MRSA/MSSA living in the nose has been shown to significantly reduce staphylococcus aureus infection. Historically, this treatment has been commonly used but growing mupirocin resistance is developing, and alternatives are being sought (Phillips et al., 2014). Currently, there are studies showing that a swab of povidone-iodine to the nares immediately before surgery is just as effective as the mupirocin ointment and compliance with treatment is improved since it is administered at the hospital before surgery (Phillips et al.). This treatment is being implemented at various hospitals around the country. The purpose of this project was to determine the effectiveness of nasal mupirocin vs. nasal povidone-iodine administered pre-operatively for the decolonization of MRSA/MSSA in the nares to prevent the incidence of post-operative infection.

Next, the literature review will be presented.

Literature Review

Post-Operative Surgical Site Infections: Definition and Incidence

The skin is the largest organ of the body and the first line of defense against infections. If the skin is breached either surgically or by trauma, the body is at risk for infection (Surahio, Talpur, Memon, Junejo, & Laghari, 2017). Surgical site infection (SSI) remains a substantial cause of morbidity and mortality as well as the cause of prolonged hospitalizations. Surgical site infections are the second most common type of adverse event that occur in hospitalized patients after surgery and are one of the most common surgical complications (Cheng, Li, Kong, Wang, Ye, & Xia, 2015).

The term ‘surgical site infection’ was developed by the Centers for Disease Control (CDC) in 1992 (Owens & Stoessel, 2008). A surgical site infection is defined by the CDC as an infection that occurs after surgery in the part of the body that the surgery took place (CDC, 2019). The CDC has specific SSI criteria which include: superficial incisional surgical site infection; deep incisional surgical site infection; and organ/space surgical site infection (CDC).

A superficial incisional surgical site infection must meet the following criteria: an infection occurring within 30 days of the operative procedure; involves only the skin and subcutaneous tissue of the incision; and has purulent drainage from the incision site, organism identified by aseptically-obtained specimen, or signs and symptoms of an infection (CDC, 2019). Criteria for a deep incisional surgical site infection include: occurs within 30-90 of the operative procedure; involves deep, soft tissues of the incision; and has purulent drainage from the incision, organisms identified via a culture swab, or at

least one sign or symptom of infection (CDC). An organ/space surgical site infection must meet the following criteria: infection occurs within 30-90 days of the operative procedure; involves any part of the body deeper than the fascia/muscle layers; has purulent drainage, organisms identified via culture, or abscess or infection identified via exam; and must be an approved organ/space infection site (CDC).

Until the middle of the 19th century, most wounds became infected and those that were extensive resulted in a mortality rate of 70-80%. The rates of surgical site infections drastically improved when Ignaz Semmelweis and Joseph Lister pioneered infection control by introducing anti-septic surgery. Despite these advances, the overall incidence of healthcare associated infections (HAIs) remains high (Bandaru, Rao, Prasad, & Murty, 2012). Surgical site infections are now the costliest and most common hospital acquired infections (Anderson et al., 2014).

Surgical site infection rates are estimated at less than 1% to more than 10%, with 75% of SSI-associated deaths directly attributed to a SSI (Cheng et al., 2015). This poses a large economic and financial burden on the patient, their family, and the healthcare institution. The incidence of SSI differs greatly depending on the hospital and from one geographic location to another. While a SSI is a serious threat to patients' lives, it is also a large financial burden on patients, families, and society (Cheng et al.). Patients with SSIs are more likely to require hospital readmission, possibly to an intensive care unit and are at higher risk of death than those without infections (Owens & Stoessel, 2008).

The financial burden of an SSI is approximately twice the amount of in-patient costs for those without a SSI. A study of hospital-acquired infections in Massachusetts

estimated the cost to be between \$223,000,000 and \$275,000,000 (Stone, Kunches, & Hirschhorn, 2009). Surgical site infections are estimated to account for 3.5 to 10 billion dollars annually in health care expenditures (Anderson et. al, 2014). Surgical site infections are associated with a mortality rate of 3% with 75% of SSI-associated deaths being directly attributed to an SSI (CDC, 2019). Therefore, is imperative that SSIs are identified and diagnosed early and are treated immediately upon diagnosis. Factors responsible for surgical site infections must be identified and measures must be taken to prevent these factors from occurring to reduce morbidity, mortality, and healthcare expenses (Cheng et al., 2015).

Risk and Contributing Factors for Surgical Site Infections

There are two different categories of factors that place a person at increased risk for a SSI. These categories are patient-related (endogenous) or procedure-related (exogenous) factors (Owens & Stoessel, 2008). Patient-related factors include advanced age, nutritional status, diabetes, smoking, obesity, coexistent infection at a remote body site, colonization with micro-organisms, altered immune response, and length of preoperative hospital stay. Procedure-related factors include duration of surgical scrub, skin antisepsis, preoperative shaving, preoperative skin preparation, duration of operation, antimicrobial prophylaxis, operating room ventilation, inadequate sterilization of surgical instruments, foreign material in the surgical site, surgical drains, and surgical technique (Owens & Stoessel). Further procedure-related factors include: contamination from members of the surgical team; the operating room environment; and instruments brought into the sterile field used during the surgical procedure. The organisms

associated with exogenous factors include staphylococci and streptococci (Owens & Stoessel).

A study by Alfonso-Sanchez, Martinez, Martin-Moreno, Gonzalez, and Botia (2017) examined the risk factors influencing surgical site infections. This was a longitudinal prospective study at eight different hospitals designed to identify SSIs in all patients who underwent surgical procedures. The patient-related variables included in this study were: age; sex; number of comorbidities; diabetes; cancer; chronic obstructive pulmonary disease; hepatic cirrhosis; smoking; immunosuppression; nutrition; nasogastric tube feeding; transfusion; and length of preoperative stay. The variables associated with the operation included: antibiotic prophylaxis; depilation; American Society of Anesthesiologists (ASA) class; type of intervention; and duration of intervention.

The sample size included 18,910 patients, with 1,267 (6.7 %) developing a SSI. The most significant environmental factor risk was contamination in the operating room by bacteria and fungi. Bacteria caused 1.75% of infections ($p < 0.001$) and fungi with greater than 6 CFU caused 6.23% of infections ($p < 0.001$). Each type of surgical infection was found to have different associated risk factors. Superficial SSIs were associated with environmental factors such as contamination by fungi and bacteria, surface contamination, humidity, differential pressure, and temperature of the operating room. Whereas the factors associated with deep organ/space SSIs were associated with patient characteristics such as age, sex, transfusion, nasogastric feeding, nutrition, type of intervention, and preoperative stay.

In addition to the above stated risk factors, the virulence of the bacteria and the procedure performed are also important determinants in the development of a wound infection (Surahio et al., 2017). Most wound infections only involve the skin and subcutaneous tissue but if left untreated or not adequately treated early and aggressively, they can lead to a more serious infection or sepsis (Surahio et al.). Presenting signs of an infection usually involve increasing pain at the wound, fever, and/or discharge from the wound. They may also present with erythema and tenderness to the incision site (Surahio et al.).

Pathogens and Surgical Site Infection

While there are endogenous and exogenous related factors for developing a SSI, the most responsible pathogens originate from the patient's endogenous flora. There are many organisms related to SSIs but the most common are staphylococcus aureus, coagulase-negative staphylococci, Enterococcus spp., and Escherichia coli with an increasing number related to antibiotic-resistant organisms such as methicillin resistant staphylococcus aureus (MRSA) (Owens & Stoessel, 2008). Patients infected with MRSA have an 18% mortality rate, even among healthy patients (Loftus, Dexter, & Robinson, 2018). Recent evidence concluded that the MRSA infection rate in 2015 was no different than the 2010-2011 baseline (Kavanagh, Abusalem, & Calderon, 2017).

Cheng et al. (2015) conducted a prospective study at a tertiary care center from July 2013 to December 2014 which included 1,138 patients aged two to 92 who underwent breast, hernia, esophagus, stomach, appendix, colon, or rectal surgery. The purpose of the study was to identify risk factors for surgical site infections in a teaching

hospital. The authors found a significant association between the age of the patient and the development of a surgical site infection. Patients aged over 75 years were more likely to develop a surgical site infection (5.6%; n= 161) than those less than 75 years of age (3.0%; n= 977). They also found that there was a correlation between those patients with diabetes (odds ratio [OR] 6.400; 95% confidence interval [CI] 2.582–15.866; $p= 0.000$); and/or cancer (OR 2.427; 95% CI 1.028–5.732; $p= 0.043$); and being at an increased likelihood for developing a SSI. Patients with diabetes are more susceptible to a wound infection because of impaired neutrophil chemotaxis and phagocytosis. There was a significant difference between the diabetic and non-diabetic groups; the percentage of surgical site infections in diabetic patients was higher (14.3%; n= 77) ($p= 30.660$) than for non-diabetics (2.5%; n= 1,061) ($p= 0.000$). Patients with cancer are at increased risk for a SSI due to immunosuppression and their reduced ability to fight off infection. The percentage of patients with an SSI who had cancer was 8.4% (n= 77, $p= 10.559$) and those without cancer was 2.7% (n= 1,019, $p= 0.001$). There was a higher incidence of surgical site infection in emergency surgery cases (8.4%; n= 166) than elective surgery cases (2.5%; n= 972) because emergency procedures do not allow for stable vital signs, adequate antiseptic skin preparation, and decontamination of the colon prior to colon surgery. P-value for elective surgery was $p= .000$ (Cheng et al.).

This study isolated 20 different organisms as causing the surgical site infections, with the most prominent being *Escherichia coli*, *S. aureus*, and *Pseudomonas aeruginosa*. Based on this, it was recommended that preventive measures should be strengthened to reduce the incidence of SSI. Furthermore, this study showed that there was an increase in surgical site infection with the increasing volume of blood transfusions. The incidence of

surgical site infection was higher (10.7%; n= 56) in those who received transfusions than in those who did not (3.0%; n= 1,082) with a P-value of 0.002. The authors concluded that patients should not receive blood transfusions unless absolutely necessary (Cheng et al., 2015).

Surahio et al. (2017) performed a prospective, descriptive study to determine the frequency of SSIs in postoperative patients and to evaluate the type of organism involved in post-operative infections and its sensitivity. All patients were older than 13 years and underwent surgery and subsequently developed a surgical wound infection. The study was performed over a one-year period and included 424 patients. The results indicated that 2.25% (n= 8) of patients developed a grade I infection, 11.58% (n= 41) developed a grade II infection, 1.41% (n= 5) developed a grade III infection, and 0.28% (n= 1) developed at grade IV infection. Twenty-three of these patients underwent an appendectomy, five had inguinal hernia surgery, nine patients had a laparotomy, and eight patients underwent a cholecystectomy (Surahio et al.).

The most common organisms isolated from the infected post-operative wounds were *Escherichia coli*, which infected 24 patients and *S. aureus* which infected six patients. When an antibiotic sensitivity was performed it showed that meropenem and piperacillin/tazobactam were effective against these organisms. Meropenem was sensitive in 96.15% (n= 25) of patients with *Escherichia coli* and 87.5% (n= 7) with *S. aureus*. Piperacillin/tazobactam was sensitive in 92.30% (n= 24) of patients with *Escherichia coli* and 75% (n= 6) with *S. aureus*. Vancomycin was found to be sensitive to *S. aureus* in 87.5% (n= 7) of patients and 100% (n= 3) of patients with streptococci (Surahio et al., 2017).

The authors concluded that wound infection caused significant morbidity in 15.53% of surgical cases and had a significant economic burden on patients and hospitals. The authors noted that antibiotics should only be used after sensitivity reports have been reviewed to avoid the development and persistence of antibiotic resistant organisms. Of most importance is maintaining optimal conditions as much as possible to avoid post-operative infections and the need to use antibiotics (Surahio et al., 2017).

Prevention of Post-Operative Surgical Site Infections: The Surgical Care Improvement Project (SCIP)

As a result of the high cost of SSIs and the increased mortality rates associated with inconsistent compliance with infection prevention measures, the Surgical Infection Prevention (SIP) project was created by the Centers for Medicare & Medicaid Services (CMS) and the CDC. The SIP was created after The Joint Commission (TJC) created four core measurement areas for hospitals: acute myocardial infarction; heart failure; pneumonia; and pregnancy related conditions in 2001 (National Hospital Inpatient Quality Measures Specifications Manual [NHIQMSM], 2014). The SIP focused on seven procedures: abdominal hysterectomy; vaginal hysterectomy; hip arthroplasty; knee arthroplasty; cardiac surgery; vascular surgery; and colorectal surgery (Anderson et al., 2014). After TJC began collecting data on the four core measurement areas, they subsequently created the Surgical Care Improvement Project (SCIP) in 2006 which included seven SCIP performance and outcome measures applicable to the peri-operative period. These measures include: prophylactic antibiotic received within 1 hour prior to surgical incision; prophylactic antibiotic selection for surgical patients; prophylactic antibiotics discontinued within 24 hours after surgery; controlled postoperative blood

glucose; appropriate hair removal; urinary catheter removed on post-operative day 1 or 2; and peri-operative temperature management (NHIQMSM).

The first SCIP outcome pertains to the timing of the first administration of prophylactic antibiotics as well as the timely discontinuation of prophylactic antibiotics (NHIQMSM, 2014). The goal of prophylactic antibiotics is to establish bactericidal tissue and serum levels at the time of the skin incision (NHIQMSM). Studies have demonstrated that the administration of systemic antibiotics prophylactically before surgery decreases the incidence of wound infections by about one half. While there are great benefits to using antibiotics prophylactically, it is also important to discontinue them in a timely manner to prevent adverse effects such as the development of antibiotic resistant pathogens or *Clostridium difficile* infections (Rosenberger et al., 2011).

For the above reasons, strict guidelines have been created concerning the administration of prophylactic antibiotics prior to surgery. Primarily, the timing of the administration of the antibiotics is important (Alexander et al., 2011). It is recommended that the antibiotics are administered within the first two hours before surgery, depending on the half-life of the medication. For antibiotics with a short half-life, such as the cephalosporins, they should be administered 30 minutes before the incision to ensure the most effectiveness. Longer acting antibiotics such as vancomycin and the fluoroquinolones should be administered one to two hours before the incision is made. Re-dosing is necessary in the use of short acting antibiotics and should be administered every three hours after the incision is made with the dosage adjusted for large body size (Alexander et al.).

The second SCIP outcome pertains to the choice of antibiotics which is made based on the half-life of the medication and the length of the surgery as well as the surgery being considered clean or dirty (Rosenberger et al., 2011). The goal of antibiotic prophylaxis is to be cost effective, safe, and a broad spectrum covering the most probable intraoperative contaminants encountered during surgery (NHIQMSM, 2014). The cephalosporins provide good early penetration into the wounds and are effective against gram-positive and gram-negative organisms commonly encountered in general surgery (Rosenberger et al., 2011). Cefazolin is the drug of choice for most surgeries, with vancomycin being the second choice but not preferred due to its' potential for antibiotic resistance (NHIQMSM). The longer acting antibiotics such as vancomycin and fluoroquinolones provide lasting penetration for longer surgical cases (Alexander et al., 2011). Cefazolin is the drug of choice for most surgeries including: burns; general surgery; genitourinary; hepatobiliary; oral/maxillofacial; orthopedic spine surgery; obstetrics; gynecology; and plastic surgery. Cefazolin plus vancomycin is recommended for cardiac surgery, neurosurgery, orthopedic, and vascular surgery (Alexander et al.).

The third SCIP outcome recommends the discontinuation of prophylactic antibiotics within 24 hours; no benefit had been noted beyond 24 hours post-operatively. Peri-operative antibiotics do not sterilize tissues but reduce the bacterial burden to an amount that may be controlled by the patient's own defenses (Alexander et al., 2011). The continuation of antibiotics beyond 24 hours leads to a risk of drug resistance and secondary infections such as *Clostridium difficile* and has not shown any benefit in SSI reduction beyond a single dose of antibiotics (Rosenberger et al., 2011).

Surgical Care Improvement Project outcome number four concerns euglycemia in the prevention of SSIs, most notably for cardiac surgery patients (NHIQMSM, 2014). Of most importance are post-operative days one and two, during which maintaining appropriate glucose concentrations reduces the incidence of SSIs (Rosenberger et al., 2011). It has been well documented that patients with diabetes who undergo surgery have an increased incidence of complications which include poor wound healing, wound infections, cardiac compromise, and death (Alexander et al., 2011). The adverse effects of poor glucose control are vast and include: disturbances of microvascular responses; inhibition of complement function; increases in pro-inflammatory cytokine levels; inhibition of chemotaxins; impaired phagocytosis and intracellular killing; disturbances in reactive oxygen species; decrease in T and B cell responses; and increased apoptosis and oxidative stress in the lymphocytes. It is recommended to closely monitor the blood glucose to less than 180mg/dL for 18-24 hours after anesthesia end time (NHIQMSM).

Surgical Care Improvement Project outcome number six relates to the method of hair removal. The accepted methods of hair removal include no hair removal or using clippers or depilatory (NHIQMSM, 2014). In a prospective study by Ko, Lazenby, Zelano, and Isom (1992) the effects of hair removal methods were investigated on suppurative mediastinitis after cardiopulmonary bypass operations on 1,980 consecutive adult patients over a two-year period. Each group was randomized to manual shaving versus electrical clipping of hair before the surgical incision. The infection rate was significantly higher in the manually shaven group (13/990) versus the electrically clipped group (4/990). It is recommended to electrically clip hair to avoid skin damage associated with shaving hair with a razor (Alexander et al., 2011).

The extensively studied topic of prevention of catheter associated urinary tract infections (CAUTI) are the focus of outcome nine from SCIP. It is recommended that the urinary catheter is removed on postoperative day one or postoperative day two (NHIQMSM, 2014). Studies have found that the likelihood of developing a CAUTI directly correlates with the duration of catheter use (Rosenberger et al., 2011). Patients with catheters in place for longer than two days post-operatively have been found to have twice the chance of developing a urinary tract infection (UTI) than those who have a catheter for less than two days (Rosenberger et al.).

The final outcome, SCIP number 10, pertains to the maintenance of normothermia peri-operatively to reduce SSIs. Normothermia during surgery, according to SCIP, is defined as at least one body temperature equal to or greater than 96.8 degrees Fahrenheit recorded within 15 minutes before the surgery end time (NHIQMSM, 2014). Studies have indicated that maintaining normothermia during surgery was associated with lower rates of SSIs. It is hypothesized that hypothermia leads to peripheral vasoconstriction and impaired immune function which leads to higher rates of SSIs (Rosenberger et al., 2011). Hypothermia also contributes to: increased blood loss and transfusion requirements; prolonged anesthesia recovery; prolonged hospitalization; increased morbid myocardial events; and increased wound infections (Alexander et al., 2011). It has been shown in animal studies that an increase in the core temperature that occurs during bacterial infections is essential for optimal antimicrobial host defense (Alexander et al.).

A study by Yi, Liang, Song, Xia, and Huang (2018) was conducted to evaluate if active warming practices during surgery reduced bleeding in patients undergoing major open operations. This was a prospective, parallel two-arm randomized controlled trial

that compared active warming and passive warming practices amongst unilateral total hip replacement surgeries and open thoracic operations. The passive warming group included those covered with un-warmed cotton blankets from pre-operative holding area to operating room to post-anesthesia care unit. The active warming group members were covered with forced-air blankets connected to a warming unit during the pre-operative holding area and in the operating room. The incidence of hypothermia (<36 degrees Celsius) during surgery was 0% in the active warming group whereas the incidence was 71.8% in the passive warming group. Most importantly, the study found that the combined volume of intraoperative blood loss for the two operations was significantly less in the active warming group (464ml) than the passive warming group (682 ml). Therefore, it was concluded that it would be beneficial to implement active warming protocols to reduce SSIs (Yi et al.)

Other Surgical Site Infection Prevention Guidelines

Further recommendations for prevention of SSIs in addition to the SCIP outcomes have been identified by the CDC. The CDC guidelines for prevention of SSIs were created in 1999 and updated in 2017. The most current CDC guidelines for the prevention of SSIs include: showering or bathing with an antiseptic agent at least the night before surgery; antimicrobial prophylaxis administered when indicated and timed appropriately; skin preparation in the operating room with an alcohol based agent; no additional prophylactic antimicrobials administered after the surgical incision is closed; maintaining blood glucose less than 200mg/dL; maintaining normothermia; oxygen administration during surgery; and refraining from the transfusion of blood products (Berrios-Torres, Umscheid, & Bratzler, 2017). Other SSI prevention guidelines include techniques for air

handling, cleaning of environmental surfaces, sterilization techniques, activities of surgical team members, surgical attire, drapes, and asepsis (Alexander et al., 2011). In regard to air handling, microbes in the air are a major source of pathogens for causing wound infections and high efficiency filters provide the best way to filter the air. Another big source of contamination in the operating room is glove perforation which significantly increases the risk of infection. Double gloving has been shown to greatly decrease glove perforation (Alexander et al.).

Post-operative infections are frequently caused by organisms that are already on the skin, most commonly *S. aureus*. In the years before 1970, studies showed that bathing with hexachlorophene before surgery reduced the risk of wound infections. This practice was later switched to chlorhexidine baths because they provide better long-term suppression of organisms; this is now the agent of choice (Alexander et al., 2011). According to the CDC guidelines from 2014, patients are advised to shower or bathe with soap or an antiseptic agent at least the night before the operative day. This is classified as a category IB strong recommendation (Berrios-Torres et al., 2017). Showering with chlorhexidine pre-operatively has been shown to be more effective in reducing the number of organisms on the skin than iodine or soap and water. People who are nasal carriers of *S. aureus* or MRSA are likely to have other body sites that are also contaminated with the same organisms. Thus, they are at risk for endogenous transmission of *S. aureus* infections (Alexander et al.).

Preventing post-operative infections does not depend on just one variable. It is a complex process and it varies based on the patient, the surgical environment, and the surgical technique. Measures need to be taken to address each of these areas. Important

guidelines to follow in the prevention of post-operative infections are: correct timing of prophylactic antibiotics; intraoperative normothermia; avoiding urinary tract catheterization; adhering to basic hand hygiene; and more recently to decolonize the skin and nares with chlorhexidine (Andersson, Bergh, Karlsson, Eriksson, & Nilsson, 2012).

Staphylococcus Aureus and Surgical Site Infections

Staphylococcus aureus. *Staphylococcus aureus* is a gram-positive bacteria that can cause a variety of infections ranging from skin and soft tissue infections to life-threatening blood infections and sepsis (Hogan et al., 2016). It is a human skin and mucosa commensal and is a frequent cause of serious infections with a high morbidity and mortality as well as increased healthcare-associated costs (Sakr et al., 2018). *Staphylococcus aureus* is identified as a bacteria that commonly causes opportunistic infections from skin infections to toxic shock syndrome, endocarditis, pneumonia, and sepsis. Due to this, it is now recognized as a common cause of hospital- acquired and community- acquired infections (Waryah et al., 2016). *Staphylococcus aureus* is dispersed from a provider's hands and patient skin surfaces and is then capable of contaminating aerosolized particles, equipment, and tools such as laryngoscope blades, laryngoscope handles, anesthesia machines, and ventilators. It has been shown to survive on surfaces for up to 360 days (Loftus, Dexter, & Robinson, 2018).

Staphylococcus aureus develops due to several virulence factors which include biofilm formation, antibiotic resistance, and the production of a wide array of toxins (Waryah et al., 2016). Primarily, a biofilm, which is a congregation of microorganisms residing in a protective extracellular matrix, forms around the bacteria (Waryah et al.).

This is followed by the second stage, which involves colonization of the bacteria by means of adherence factors or adhesions which aid in attachment of the bacteria to the host surface. This is done by a microbial surface component which recognizes adhesive matrix molecules. There are over 20 different adhesive matrix molecules that can express *S. aureus* (Waryah et al.).

Once the bacteria adhere to the host surface, the biofilm is strengthened further by an intracellular adhesion encoded by a cell surface polysaccharide and an antigen. Furthermore, *S. aureus* produces a variety of endotoxins which contribute to host tissue membrane disruption and provide nutrients that are essential to bacterial cell growth (Waryah et al., 2016). Given these factors, combined with the over use of antibiotics, development of persistent antibiotic resistance has improved the ability of *S. aureus* to resist treatment with antibiotics (Waryah et al.).

Bacteria that produce biofilms contribute to greater than 80% of all infections in humans (Piechota et al., 2018). These bacteria are a primary cause of healthcare-associated infections because the biofilm show increased resistance against standard antimicrobial treatment and host immune factors and is able to colonize medical surfaces such as catheters and other devices (Piechota et al.). A study by Piechota et al. (2018) aimed to investigate the capacity of clinical strains of *S. aureus* to form biofilms. They collected *S. aureus* strains from two hospitals between 2015 and 2017 and divided them into two groups: MSSA strains and MRSA strains. There were 57 strains of MSSA and 73 strains of MRSA. All strains were evaluated for biofilm production and out of 130 strains, 99.2% were biofilm producers. Due to the high incidence of biofilm formation by both MSSA and MRSA, there is a high ability of these strains to persist in hospital

environments and increase the risk of drug resistant infections in hospitalized patients (Piechota et al.).

Staphylococcus aureus is one of the most common causes of bacteremia and currently carries a 20-40% mortality rate after an infection of up to 30 days despite medical treatment (Piechota et al., 2018). *Staphylococcus aureus* can cause opportunistic disease in the hospital as well as the community setting (Kim et al., 2018). Over the last 20 years, infections caused by *S. aureus* have become more dangerous and costly to treat due to the increasing prevalence of antibiotic resistance due to the widespread use of antibiotics (Piechota et al.). *Staphylococcus aureus* initially responded well to the beta-lactam group of antibiotics, but with the evolution of MRSA being reported in such a short period of time, it is now resistant to most antibiotics except for vancomycin (Hussain, Naqvi, & Sharaz, 2019). Methicillin resistance is mediated by a penicillin-binding protein encoded by a gene that allows the organism to grow and divide in the presence of methicillin and other beta-lactam antibiotics (Sfeir et al., 2014). The primary drug resistant strains of *S. aureus* are methicillin-sensitive *staphylococcus aureus* (MSSA) and methicillin-resistant *staphylococcus aureus* (MRSA). Methicillin sensitive *staphylococcus aureus* causes approximately half of all healthcare associated *S. aureus* infections (Kourtis et al., 2019).

Methicillin Resistant *Staphylococcus Aureus*. Methicillin resistant *staphylococcus aureus* infections first began being reported in 1950 and infection rates have increased dramatically in recent decades, reaching up to 50%, with 30% of patients infected with MRSA dying within 30 days (Al-Tamimi et al., 2018). Hospitals are the main source of MRSA outbreaks; this is due to a variety of factors including prolonged

hospital stay, widespread use of antibiotics, nursing home exposure, immune suppression, improper antibiotic dosage, indwelling catheterization, invasive medical devices, drug abusers, and unsterilized instrumentations (Hussain et al., 2019). The prevalence of MRSA has increased in the United States over the past 10 years from 32.7% in 1998 to 53.8% in 2007, with MRSA related hospitalizations doubled (Hussain et al.). Methicillin resistant staphylococcus aureus is associated with severe and prolonged infections, with invasive MRSA infections reported to have an 18% mortality rate among healthy patients in the community (Loftus et al., 2018). Methicillin resistant staphylococcus aureus has been shown to be more likely than MSSA to be implicated in intraoperative clonal transmission (Loftus et al.).

In a study by Hussain et al. (2019), the authors aimed to identify the prevalence of MRSA in a tertiary care hospital. They performed a cross sectional study over a six-month period during which 100 clinical specimens of pus and wounds were collected. The findings showed that out of the 100 clinical specimens, 65 of them showed bacterial growth. Of these 65 samples, 27 were gram positive cocci, and 38 were gram negative rods. Of the 27 gram positive cocci, 21 were *S. aureus*. Of the 21 *S. aureus* samples, 14 were MRSA and seven were MSSA. It is important to identify this nosocomial pathogen early to prevent its' dissemination and life-threatening complications.

While the MRSA rate is higher in hospitalized patients, there is growing frequency of community acquired MRSA (Al-Tamimi et al., 2018). Community acquired MRSA infections were first identified in the 1990s, which caused mostly soft tissue infections (Kourtis et al., 2019). It was first reported among injection drug users and is now the most frequent cause of skin and soft tissue infections (Sfeir et al., 2014).

Community acquired MRSA can also cause bacteremia, infective endocarditis, pneumonia, bone and joint infections, (Kim et al., 2018) necrotizing fasciitis, wound infections, otitis media, otitis externa, osteomyelitis, urinary tract infections, sepsis, and necrotizing pneumonia (Sfeir et al.).

Community outbreaks have been reported in native and aboriginal communities, sports teams, child-care centers, military personnel, men who have sex with men, and prison inmates and guards (Sfeir et al., 2014). Risk factors for development of a community MRSA infection include: skin trauma; cosmetic body shaving; incarceration; sharing of unclean equipment between users; and physical contact with others who have MRSA colonization or a MRSA infection (Sfeir et al.). Community acquired MRSA is now seen with increasing frequency in the hospital setting due to patients who acquire MRSA in the community becoming hospitalized and subsequently transmitting the community strains to other hospitalized patients (Sfeir et al.).

Staphylococcus aureus is a common bacteria in both the hospital and community setting, posing serious risks to the population. With the evolution of *S. aureus* into the resistant strains of MRSA and MSSA, it is necessary to be even more vigilant to identify and treat these infections early and prevent the transmission of disease. Since *S. aureus* can colonize a person's skin and mucus membranes, nasal colonization with *S. aureus* has been at the forefront of recent research in the prevention of post-operative infections.

Nasal colonization with Staphylococcal Aureus

In the United States, more than 40 million people undergo surgery each year, with more than 20 percent of them acquiring a nosocomial infection during the post-operative

period. Post-operative infections at the surgical site are the third most common nosocomial infection and complicate 1-10% of operations. *Staphylococcus aureus* is the cause of 25% of nosocomial infections. It colonizes the anterior nares in 25-30% of the population and those who are colonized with *S. aureus* are at higher risk for staphylococcal infections after invasive medical or surgical procedures than those who are not colonized (Perl et al., 2002). Nasal colonization is a preoperative risk factor for MRSA and MSSA infections as it can spread from the anterior nares to other areas of the skin and contaminate the incision during surgery (Sakr et al., 2018). Studies have shown that around 80% of strains that have caused a staphylococcal infection at the surgical site have the same molecular identity as the *S. aureus* strain in the nares of the infected patient (Saker et al.).

Colonization of *S. aureus* in the nares has been shown to play a role in the pathogenesis of *S. aureus* infections in patients undergoing surgery, dialysis, and intensive care unit patients (Sakr et al. 2018). *Staphylococcus aureus* establishes solid interactions with nasal epithelial cells via various proteins and many cell surface components. The anterior nares are lined by a stratified, keratinized nonciliated squamous epithelium, where the rest of the nasal cavity is lined with a ciliated columnar epithelium. The outermost layer of the anterior nares is the stratum corneum, which contains keratinocytes that express proteins. These proteins are able to interact with staphylococcal surface proteins which favor nasal colonization (Sakr et al.). Colonization depends on the host's defenses due to underlying conditions or diseases. Healthy hosts have lower rates of colonization whereas those with conditions such as human

immunodeficiency virus (HIV), obesity, diabetes, and dialysis patients have been found to have higher rates of nasal colonization (Sakr et al.).

Staphylococcus aureus can be found in various sites on the body including the skin, rectum, vagina, gastrointestinal tract, and axilla, with the main reservoir being the anterior nares (Sakr et al., 2018). When in contact with the nasal mucosa, *S. aureus* then interacts with epithelial cell ligands. Once the bacteria overcome the host's defenses, it can propagate into the anterior nares and the host becomes a *S. aureus* nasal carrier. Nasal carriage can begin within the first days of life, with 90% of *S. aureus* strains found to be identical to the maternal nasal strain (Sakr et al.). After birth, the hands are the main source of infection, with transmission occurring from surfaces to the hands and then the nose.

In a study by Loftus et al. (2018), the researchers aimed to identify the source of transmission for *S. aureus* in the operating room of 274 case pairs of three academic medical centers in the United States. This study involved the first and second cases of the day. Before the first case of the day, samples were taken from the bacterial reservoirs. Bacterial reservoirs for this study included: the anesthesia providers' hands before, during, and after patient care; the adjustable pressure-limiting valve and agent dial of the anesthesia machine; the patients' nasopharynx and axilla; other providers' hands present in the operating room; and the internal lumen of the patients' intravenous stopcock set. These reservoir sites were then sampled after completion of the surgery as well. Before the first case of the day, each environmental site was decontaminated but the sites were not additionally decontaminated before the second case except for usual routine cleaning procedures to assess for the efficacy of routine cleaning procedures. The results of the

study showed that there were 22 *S. aureus* strains isolated from intraoperative reservoirs. Of the 173 *S. aureus* isolates that were collected, the source of the infection was linked to either the patient, the residents' hand, or unknown. The conclusion of this study indicated that preoperative patient decolonization and perioperative hand hygiene infection control measures need to be improved upon.

Airborne transmission is another possible route, as the risk of disseminating *S. aureus* during a viral upper respiratory infection increases the risk of infection outbreaks (Sakr et al., 2018). Air transmission is of particular concern in the operating room where it is considered a continuous medium that is able to spread infection to various sites due to the setline of aerosolized particles (Loftus et al., 2018). Healthcare workers who are asymptomatic carriers of *S. aureus* can also be the source of MRSA outbreaks as well as mobile phones contaminated with *S. aureus* (Sakr et al.).

Healthcare workers who are asymptomatic carriers of *S. aureus* are responsible for some of the transmission of *S. aureus* infections in hospitals. This was studied by Al-Tamimi et al. (2018) amongst medical students. The authors performed a cross-sectional study which included 290 medical students from the first to fourth year at a medical university. Nasal swabs were collected from each participant under sterile conditions. Out of the 290 nasal swabs, 66 of them (22.7%) were identified as being infected with *S. aureus*. Out of these 66 swabs, 54 of them (18.6%) were identified as MSSA and 12 (4.1%) were identified as MRSA. Potential risk factors for nasal colonization found in this study included the male sex and chronic illnesses. The study concluded that nasal colonization with *S. aureus* plays an essential role in transmission of infections and those

colonized in their nares with MRSA possess a higher risk for transmission of nosocomial infections.

Another study that evaluated the prevalence of MSSA and MRSA among healthcare workers was performed by Hogan et al. (2016). These researchers performed a cross sectional study where they screened nasal swabs from 863 healthcare workers and 685 students for *S. aureus*. Of these 1548 samples, 171 were isolated as having *S. aureus*. The prevalence of *S. aureus* colonization in the healthcare worker group was 10.4% and 11.4% in the student group. Methicillin resistant staphylococcus aureus colonization was higher in the healthcare group (1.5% versus 0.9%). Nasal colonization was higher among women than men and those greater than the age of 25. Colonization of *S. aureus* was slightly higher in healthcare workers that reported direct contact with patients.

Treatment of Nasal Colonization with Mupirocin

Staphylococcus aureus is able to establish colonization in the nasal mucosa, which begins the pathogenesis of disease. Approximately 20% of people are persistently colonized with *S. aureus* and 60% are intermittently colonized in their nares. When patients are hospitalized or develop immune compromise and are colonized with *S. aureus* in their nares, they are at increased risk of developing a bloodstream infection (Uchiyama et al., 2019). Therefore, at-risk patients are screened for MRSA colonization and are decolonized, especially before surgery. Mupirocin has typically been used in the past but due to increased resistance to the drug (approaching 30% in some clinical populations) and repeated applications required, other alternatives are being sought (Uchiyama et al.).

The most common treatment for eradication of *S. aureus* in the nares is with the topical treatment of 2% mupirocin calcium ointment (bactroban) which decolonizes the anterior nares and aids in preventing post-operative infections; this has been supported by several studies which have reported lower rates of surgical site infections after treatment pre-operatively with mupirocin (Perl et al., 2002). Perl et al. (2002) conducted a clinical trial to determine whether the application of intranasal mupirocin ointment pre-operatively would decrease the rate of *S. aureus* infections at surgical sites. The study was a randomized, double-blind, placebo-controlled clinical trial at two hospitals in Iowa over a period of three years. The study subjects were treated with mupirocin ointment or placebo to the anterior nares twice daily for up to five days before surgery and monitored for 30 days postop to determine whether they acquired *S. aureus*.

The results of the study showed that the rate of infection at the surgical site was 7.9% (n= 1,933) in the mupirocin group and 8.5% (n= 1,931) in the placebo group. The study also found that the risk of nosocomial infection with *S. aureus* at any site among patients with nasal colonization of *S. aureus* was significantly lower among those who received mupirocin (12.8%; n= 444) than those who received the placebo (16.1%; n= 447) (p= 0.02). Of the participants (n= 129) with nosocomial infections who had nasal carriage of *S. aureus*, wound cultures were obtained from 107 (43 in the mupirocin group and 64 in the placebo group). Of those who were *S. aureus* carriers, 17 carriers who received mupirocin developed nosocomial *S. aureus* infections whereas 34 carriers developed a nosocomial infection in the placebo group. Among those who received the placebo and were *S. aureus* carriers, there was a 4.5 times higher chance of developing a SSI than non-carriers (Perl et al., 2002).

While mupirocin has been commonly used to treat *S. aureus* colonized in the nares before surgery, there have been reports of emerging mupirocin resistance as well as decreased rates of compliance due to the medication needing to be applied by the patient for 5 days before surgery (Phillips et al., 2014). In a study by Ramos et al. (2011), it was noted via a patient survey that while 94% of patients used the chlorhexidine soap as prescribed before surgery, only 86% actually applied the mupirocin ointment to their nares as prescribed and 8% of patients found that it was hard or very hard to purchase the mupirocin ointment due to its' cost. Therefore, other alternatives to mupirocin have been investigated and intranasal povidone-iodine has shown promising results.

Nasal colonization depends on many factors including host defenses, environmental, bacterial, and exposure to *S. aureus*, but once a person is colonized it places them at increased risk for subsequent infections. It is recommended that patients undergo decolonization of the nares prior to surgery to prevent post-operative surgical site infections. Treatment with mupirocin to both nares for five days before surgery has been recommended but current studies are being performed evaluating the efficacy of intranasal povidone-iodine administered in the hours before surgery as another alternative. The effectiveness of intranasal mupirocin versus intranasal povidone-iodine are compared in this integrative review.

Treatment of Nasal Colonization with Povidone-Iodine

Povidone-iodine was discovered in 1812 by a chemist from France and was first documented for use as an antiseptic on wounds in 1839 (Zinn et al., 2010). Iodine is currently available in the form of povidone-iodine (PVP-I) or cadexomer iodine. As

cadexomer iodine has been reported to cause transient patient discomfort with its' use, PVP-I is frequently used (Campbell & Campbell, 2013). Iodine is the bactericidal component of PVP-I combined with polyvinylpyrrolidone (povidone) which is a synthetic polymer and is commonly found in a 10% solution in water, yielding 1% available iodine (Burks, 1998).

Iodine is an essential element and has a long history as a disinfectant and antibacterial sterilizing agent, especially in surgical skin preparation. The 3M company began marketing a skin and nasal antiseptic preparation (SNP) based on povidone-iodine (PVP-I) as an alternative to mupirocin in 2010 (Uchiyama et al., 2019). Povidone-iodine has rapid in vitro activity and the duration of the effect on the skin can last 12-14 hours due to a process called back diffusion (Anderson et al., 2015). In a study by Anderson et. al (2015), it was concluded that the 3M company's SNP achieves a significant reduction in the resident *S. aureus* from the anterior nares of human test subjects.

Iodine demonstrates rapid and broad-spectrum bactericidal activity within 10-20 seconds and there is no evidence that bacteria can develop resistance to iodine since it reacts rapidly with double-bonds, amino groups, and solphydral groups, resulting in simultaneous action against multiple molecular targets to cause cell death (Uchiyama et al., 2019). Povidone-iodine works by releasing free iodine which binds to the bacteria (Zinn et al., 2010). Povidone-iodine is rapidly microbicidal with multiple mechanisms of action which include binding to proteins, nucleotides, and cell membrane fatty acids which all disrupt cell function (Campbell & Campbell, 2013). This mechanism of action coupled with the rapid action of iodine give iodine its' ability to avoid drug resistance. Iodine has excellent microbicidal activity because it has a very broad spectrum of action.

It is the only topical antimicrobial agent that is effective against both gram-positive and gram-negative organisms, spores, amoebic cysts, viruses, fungi, protozoa, yeasts, and MRSA (Campbell & Campbell).

Studies involving the use of povidone-iodine to treat nasal colonization with *S. aureus* in the prevention of SSIs have emerged due to growing mupirocin resistance, cost of the drug, and improper usage by patients. In the search for alternatives to mupirocin, povidone-iodine has gained increased interest because it is a broad-spectrum antiseptic that is suitable for the suppression of *S. aureus* (Phillips et al., 2014). In contrast to mupirocin ointment intranasally to eradicate *S. aureus* in the nares before surgery, applying povidone-iodine to the nares one time just prior to surgery is intended to transiently suppress *S. aureus* in the nares during surgery (Phillips et al.). Therefore, with the use of povidone-iodine, user error is removed as the nurse would be applying the solution before surgery; this would also potentially lower cost.

Next, the theoretical framework used to guide this integrative review will be presented.

Theoretical Framework

The purpose of a theoretical framework is to create boundaries for a project, create a structure within the boundaries, define concepts used, increase efficiency in the project, create organization, and provide consistency (Bonnell & Smith, 2018). The theoretical framework chosen for this integrative review is the Iowa Model of Evidence-Based Practice. This model was first developed in 1994 by a team of nurses from the University of Iowa Hospitals and Clinics (UIHC) to guide clinicians in evaluating and infusing research findings into patient care (Iowa Model Collaborative, 2017). The Iowa Model was revised in 2001 to reflect the evolution into evidence-based practice (EBP) involving multiple levels of evidence (Iowa Model Collaborative, 2017). It was again revised in 2015 to reflect even more dramatic changes in health care. The Iowa Model version from 2015 will be used for the purposes of this integrative review.

The Iowa Model is comprised of a flow chart (Figure 1) which consists of: identify triggering issues/opportunities; assemble, appraise, and synthesize body of evidence; design and pilot the practice change; and integrate and sustain the practice change. Underneath each phase there are sub-headings with further considerations. After each phase, there is a question to determine if the user should continue on with the question or consider another opportunity. An important part of the Iowa Model worth noting comes after the identify triggering issues/opportunities phase, where the question or purpose of the investigation must be stated. This is used to conduct a literature search for research studies that pertain to the question at hand (Brown, 2014).

The Iowa Model Revised: Evidence-Based Practice to Promote Excellence in Health Care

Figure 1. The Iowa Model flow chart. This figure illustrates the 4 phases to use evidence-based practice to promote health care.

Next, the methods to be used in this integrative review will be presented.

Methods

Purpose of Study/Clinical Question

The purpose of this project was to conduct an integrative review to examine the effectiveness of nasal mupirocin vs. nasal povidone-iodine administered pre-operatively for the treatment of a MRSA/MSSA infection to prevent the incidence of post-operative infection.

The question posed was: What is the effectiveness of mupirocin vs. iodine administered nasally pre-operatively at preventing post-operative infections?

Outcomes Examined

The outcomes examined were development of a post-operative infection, decolonization of MSSA/MRSA, and cost effectiveness of povidone-iodine.

Inclusion and Exclusion Criteria

Inclusion criteria included: adult patients 18 years and older; colonization with MRSA/MSSA in the nares pre-operatively; nasal colonization confirmed via nasal swab; inpatient and outpatient surgical patients; measurement of post-operative infection as an outcome; measurement of MRSA/MSSA rates as an outcome; quantitative studies and literature reviews comparing the effectiveness of mupirocin versus povidone-iodine published from January 2014-January 2020; quantitative studies and literature reviews related to the effectiveness of mupirocin; and quantitative studies and literature reviews related to the effectiveness of povidone-iodine. Exclusion criteria included: quantitative

studies and literature reviews before January 2014 and those not meeting the inclusion criteria were excluded.

Search Strategy

The databases searched included PubMed, CINAHL, and Google Scholar. The terms searched included: Nasal eradication with mupirocin and prevention of post-operative infection and nasal eradication with povidone iodine and prevention of post-operative infection. The selection of research articles was guided by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram (Figure 2). The flow diagram, as illustrated on the next page, was used to document the studies and the process of selection to be used in the review. The flow diagram identified the literature search, the number of records identified, the number of records screened and excluded, and the number of studies included (Moher et al., 2009). The records that did not meet the inclusion criteria and duplicates were excluded. The final result was a total of 6 articles included in this integrative review.

Figure 2. The PRISMA flow diagram. This figure illustrates the number of records identified, included, excluded, and reasons for exclusion.

Data Collection and Appraisal

The tools used for the data collection and critical appraisal of the included studies were Polit and Beck's Guide to an Overall Critique of Qualitative Research, Quantitative Research Report, and Literature Reviews (2017). For the purposes of this integrative

review the quantitative and literature review guidelines were utilized. Tables reflecting the critiques of the quantitative research studies are presented in Appendix A.

The quantitative review guidelines consisted of six major headings including the title, abstract, introduction, method, discussion, and general issues. The introduction included sub-headings such as statement of the problem, hypotheses, literature review, and theoretical framework. The method section included protection of human rights, research design, population and sample, data collection and measurement, procedures, data analysis, and findings. The discussion section included interpretation of the findings and implication/recommendations. General issues addressed presentation, researcher credibility, and summary assessment. Each sub-heading had one or more critiquing questions associated with it.

The literature review guideline consisted of seven critiquing questions: how thorough the review was; the sources the review relied on; if key studies were critically appraised; the organization of the review; objectiveness of the review; if the review was part of a research report for a new study; and if the article drew reasonable conclusions about practice implications.

Data collection was performed using a data collection tool created specifically for this review (Table 1). This collection tool provides detailed, pertinent information on each research study involved in this integrative review and includes the purpose, methods, and results of each research article. Tables reflecting the results of the data collection tool can be found in Appendix B.

Table 1

Data Collection Tool: Background Information

Purpose	Methods	Results

Cross Study Analysis

Cross study analysis was performed using descriptive data synthesis. This was attained in a narrative form to summarize and compare each study as well as to identify common findings. A table was created to help visualize the findings across the studies (Table 2). Throughout the cross study analysis, patterns and themes were identified as well as commonalities and differences. Tables reflecting the cross-study analysis can be found in Appendix C.

Table 2

Data Collection Tool: Cross Study Analysis

Author	
Key Findings	
Recommendations	

Next, the results of this integrative review will be presented.

Results

The term “nasal eradication with mupirocin and prevention of post-operative infections” was searched and yielded 6,820 results on Google Scholar, Cinahl yielded 453 results, and PubMed yielded 0 results. After narrowing date of publication to between the years of 2014-2020, Google Scholar yielded 3,220 results and Cinahl yielded 148 results. After filtering out articles that were not available in full text, four results met the inclusion criteria on Google Scholar and one from CINAHL.

When the term, “nasal eradication with povidone-iodine and prevention of post-operative infection” was searched and yielded 7,190 results on Google Scholar, 1,047 results on CINAHL, and 0 results on PubMed. After narrowing the date of publication to between the years of 2014-2020, Google Scholar yielded 3,190 and CINAHL yielded 398 results. After filtering out articles that were not available in full text, six articles met the inclusion criteria on Google Scholar and two articles on CINAHL. After filtering out duplicate articles from each search, a total of six articles were included in this integrative review. The PRISMA flow diagram on the next page illustrates the search strategy.

Figure 3. PRISMA flow diagram. This figure illustrates the process for selecting the studies to be used in a systematic review.

Results for this integrative review will be presented in order from earliest date of research. Tables consisting of critiques from each study reviewed are presented in Appendix A.

Phillips et al. (2014) (Appendix A-1) conducted an investigator initiated, prospective, open label, randomized trial comparing SSI after arthroplasty or spine fusion in patients receiving topical 2% chlorhexidine gluconate wipes with either twice daily application of mupirocin 2% ointment for five consecutive days prior to surgery or two 30 second applications of povidone iodine 5% solution into each nostril within two hours of surgical incision. The study end point was SSI within three months following surgery caused by any pathogen.

This study was performed from March 2011 through March 2012 and subjects were at least 18 years old who were having primary or revision arthroplasty and spine fusion surgery. Subjects were stratified by arthroplasty or spine fusion surgery and then randomized equally to either the mupirocin or povidone iodine treatment groups. The two intervention groups received two applications of six chlorhexidine wipes to the skin from the chin to toes, received appropriate perioperative antimicrobial prophylaxis, and either 7-10 applications of mupirocin to the nares over the five days prior to surgery or two applications of povidone iodine to each nostril within two hours of surgical incision.

The findings are presented in Appendix B-1. Phillips et al. (2014) conducted a 12-month enrollment period and 1,874 of the 1,903 patients assessed were enrolled and randomized, 177 of which did not receive the study intervention due to the surgery being cancelled or the actual surgical procedure performed was not eligible for inclusion in this

study. An SSI caused by any deep surgical site infection developed in 14 of 855 surgeries in the mupirocin group (n= 855) and six of 842 surgeries in the povidone iodine group (n= 842) (p= 0.1). In the per protocol analysis, 13 of 763 subjects in the mupirocin group developed any deep surgical site infection (n= 763), compared to 5 of 776 subjects in the povidone iodine group (n= 776) (p= 0.06). Deep surgical site infection caused by *S. aureus* occurred in 5 of 855 subjects in the mupirocin group (n= 855) and 1 of 842 subjects in the povidone iodine group (n= 842) (p= 0.2) in the intent to treat group. The intent to treat group included those who were enrolled and met eligibility requirements for the study but did not complete the assigned study regimen. In the per protocol group, 5 of 763 subjects in the mupirocin group developed a deep surgical site infection caused by *S. aureus* (n= 763) and 0 of 776 subjects in the povidone iodine group (n= 776) (p= 0.03).

The authors concluded that povidone iodine is preferred due to its' ease of use and high compliance rates. Povidone iodine is given just prior to surgery by a medical professional, therefore reducing the risk for patient error. Mupirocin, on the other hand, is administered at home by the patient prior to surgery, leaving room for patient error or noncompliance. At \$20 per application, povidone iodine provides more value, defined as quality of outcomes divided by cost, as compared to mupirocin at \$130 per course.

Anderson et al. (2015) (Appendix A-2) conducted a quantitative study to examine the efficacy of nasal povidone iodine treatment in explant models and human subjects in the decolonization of nasal *S. aureus*. Baseline nasal swab samples of the anterior nares were taken from healthy human subjects (n= 70) prior to application of povidone iodine or saline. Only subjects with baseline levels of $>5 \times 10^3$ CFU/swab were included in

this study. Subjects either applied the povidone iodine solution to their nostrils or the 0.9% saline control to the nostril for 30 seconds each, followed by an immediate repeat application. Swabs were repeated one hour, six hours, and 12 hours after application of the povidone iodine or 0.9% saline control.

Anderson et al. (2015) (Appendix B-2) compared the antimicrobial effect of povidone iodine to bactroban nasal on 10 Mup MRSA isolates, both high-level resistance and low-level resistance, of ex vivo porcine vaginal mucosa was evaluated. Statistical results showed that the povidone iodine treated explants had significantly less MRSA bacteria than untreated controls and bactroban nasal treated explants (1.63 ± 0.44 versus 5.30 ± 0.30 and 5.71 ± 0.57), respectively. The low-level MRSA isolates ($n=4$) at 1-hour post application yielded a p-value of <0.05 . The high-level MRSA isolates ($n=6$) at 1-hour post application yielded a p-value of <0.05 . These values were the same at 6 and 24 hours post application.

The efficacy of povidone iodine on normal flora in the anterior nares of human subjects was also evaluated. Baseline samples ($n=70$) were obtained by swabbing the anterior nares. The anterior nares were sampled at one, six, and 12 hours following application of povidone iodine ($n=13-18$) or saline control ($n=7-9$). At all three points, *S. aureus* reduction from the baseline level in povidone iodine treated subjects was significantly greater than the saline control subjects (2.3 ± 1.68 versus 0.86 ± 0.73 at 1h; 2.79 ± 1.52 versus 0.76 ± 0.58 at 6h; and 2.37 ± 1.77 versus 0.6 ± 0.9 at 12h [$p=<0.05$]).

The authors found that when using the ex vivo porcine vaginal mucosal model, treatment with povidone iodine was bactericidal against MRSA within two hours of application. The explants treated with the 2% mupirocin ointment showed no change in the MRSA burden 14 hours post application, which is consistent with its' slow mode of action, requiring it to be applied for five consecutive days before surgery. This contributes to patient noncompliance. Furthermore, treatment with povidone iodine resulted in sustained bactericidal activity for up to eight hours after application likely as a result of increased adhesion to the mucus on the tissue surface.

The authors concluded that treatment with povidone iodine showed a significant reduction in *S. aureus* from the anterior nares of human test subjects as well as ex vivo human skin models. Therefore, due to medication non-compliance, evolving mupirocin resistance, and the need to reduce the risk of *S. aureus* SSIs, povidone iodine should be considered as an alternative to nasal mupirocin due to its rapid efficacy, broad-spectrum activity against multiple opportunistic pathogens, lack of development of antimicrobial resistance, and ease of use.

Bebko et al. (2015) (Appendix A-3) performed a prospective, retrospective review of de-identified clinical data study at the Michael E. DeBakey Veterans Affairs Medical Center (MEDVAMC). The study was performed to assess the effect of a decontamination protocol on SSIs. The protocol consisted of the application of chlorhexidine washcloths 2% and oral rinse 0.12% the night before and the morning of the day of surgery along with the intranasal povidone-iodine solution 5% once on the morning of the day of surgery for patient undergoing elective orthopedic surgery with hardware implantation.

The orthopedic service was chosen for this study because they were a high outlier with regard to SSI rates. This decontamination protocol was implemented in May 2013. Patients operated on from October 1, 2012 to April 30, 2013 were part of the control group and those operated on from May 1, 2013 to December 31, 2013 were the intervention group. All patients were monitored for 30 days postoperatively. There was a total of 344 patients in the control group and 365 patients in the intervention group.

The results (Appendix B-3) of the study revealed 3.8% or thirteen surgical site infections (n= 344) in the control group and 1.1% or four surgical site infections in the intervention group (n= 4), p-value= .02. Of the control group (n= 344), seven were superficial SSIs, five were deep SSIs, and one was an organ/space SSI. Of the intervention group (n= 365), two were superficial SSIs, two were deep SSIs, and zero were organ/space SSIs. There was a significant reduction in the number of SSIs (69.2%) (p=.02) after the decontamination protocol. Furthermore, decontamination was an independent protective factor against the development of an SSI (95% CI, 0.08-0.77) (p=.02). The data also showed a significant reduction in MRSA nasal carrier status in the intervention group compared with the control group with only 5 positive MRSA swabs in the intervention group (n= 365) and 14 positive MRSA swabs in the control group (n= 344) (p= .05).

This was a two-day protocol and there was a 100% adherence rate. The authors noted that wider implementation of a regimen without the need for *S. aureus* carrier identification and selective decolonization would allow for cost savings if implemented broadly. While this study was promising, it should be further evaluated through large-scale randomized controlled clinical trials.

Sai et al. (2015) (Appendix A-4) conducted a retrospective cohort study to evaluate the effectiveness of two decolonization protocols for newly diagnosed MRSA colonization in patients and to assess the impact of decolonization on the rate of MRSA infection. The study was performed at an 864-bed academic hospital in Brussels, Belgium. There was a MRSA surveillance and control program in place since 1990 and screening for MRSA via swabbing of the anterior nares and throat was routinely performed on patients with a prior history of MRSA colonization, who have been hospitalized or received antibiotics in the last six months, were admitted from other hospitals or long-term care facilities, or patients with wounds, skin lesions, or foreign material. The study included MRSA-positive patients from January 2006-June 2010.

The decolonization treatments consisted of the application of intranasal mupirocin 2% and washing with chlorhexidine soap or the application of intranasal povidone-iodine and washing with povidone-iodine soap, with each treatment lasting for five days. The mupirocin/chlorhexidine soap was used for uncomplicated cases and the povidone-iodine was used for complicated cases. Patients were swabbed every 48 hours and at least three successive nose and throat swabs that were negative were required before the patient could be discharged.

There were a total of 1150 patients admitted to the hospital that were colonized with MRSA over the study period but only 268 patients were eligible for the study. The findings are presented in Appendix B-4. Overall, 104 of the 268 patients were successfully decolonized and 164 patients were not. The success rate of the mupirocin/chlorhexidine intervention group was 56% and the success rate of the povidone-iodine group was 23%. It should also be noted that the use of povidone-iodine

was restricted to the complicated cases. Of the patients with >2 wound sites, 38 had successful decolonization (n= 104) and 90 had unsuccessful decolonization (n= 164). Of those who were only colonized in their nose, 30 had successful decolonization (n= 104) and only 11 had unsuccessful decolonization (n= 164).

This particular study demonstrated an overall MRSA decolonization rate that was lower than previously published studies. This was attributed to patients with less than ideal prognostic factors such as chronic wounds who are commonly excluded from similar studies. This study was also limited by the number of patients that did not qualify for the study due to a short length of hospital stay. While the success rate of decolonization was not high in this study, the authors recommended continuation of the identified strategy due to the effectiveness of decolonization on the infection rate.

Peng et al. (2018) (Appendix A-5) conducted a prospective, cross-sectional study conducted at the Department of Orthopedics at Peking Union Medical College Hospital. It was conducted between August 2015 and February 2016 for patients undergoing joint arthroplasty and spine fusion procedures requiring at least three days overnight in the hospital. The aim of the study was to assess the prevalence of MSSA/MRSA in eligible patients with a secondary aim of determining whether the current treatment protocols result in successful decolonization of MSSA/MRSA.

Patients were screened for nasally for MSSA/MRSA within 24 hours of admission by swabbing the anterior nares and then re-swabbing the day of surgery. Those who were positive for MSSA/MRSA underwent the decolonization procedure which consisted of 5% povidone-iodine nasal swabs in both nostrils twice a day for five days

prior to surgery, performed by a nurse. Patients were also instructed to take chlorhexidine gluconate baths for five days before surgery. Those who swabbed negative for MSSA/MRSA were not decolonized. Five hundred forty-five patients were screened for MRSA and MSSA and a total of 578 elective procedures were performed during that period. Of those screened, 64 were MSSA carriers and eight were MRSA carriers; therefore 72 patients were treated with 5% povidone-iodine nasal swabs for decolonization.

Findings by Peng et al. (2018) (Appendix B-5) demonstrated that the day of surgery nasal swab screening results revealed none of the patients were positive for MRSA colonization ($p = <0.000$), but 11 patients (2%) were positive for MSSA ($n=545$), which was a reduction of 92.6% from the screening before decolonization ($p = <0.001$). The eradication of MSSA colonization was 94% ($n= 545$) while the eradication of MRSA colonization was 100% ($n= 545$). The results were positive showing that this decolonization protocol was less expensive than other protocols and reduced the colonization of MSSA/MRSA in nasal carriers. The authors noted that this treatment was as effective as nasal mupirocin in successfully eradicating MRSA and significantly reducing MSSA.

Urias et al. (2018) (Appendix A-6) performed a retrospective study at a rural hospital with trauma patients undergoing orthopedic operations conducted at a medical center from October 1, 2012 through September 30, 2016. The purpose of the study was to measure the effectiveness of using CHG baths and intranasal povidone-iodine pre-operatively and its effectiveness in reducing SSIs. The trauma patients included in this study were those undergoing operative repair of lower extremity fractures using

hardware. The pre-implementation period took place from October 1, 2012 through September 30, 2014 and the decontamination protocol consisted of bathing patients with 2% CHG washcloths or Dynahex 4% CHG solution the night and morning before surgery. The implementation period took place from October 1, 2014 through September 30, 2016 and consisted of the decontamination protocol with the addition of intranasal povidone-iodine within one hour of incision.

Over the four years of this study, 1,746 trauma patients underwent a total of 1,892 orthopedic operations to repair fracture of the lower extremities using hardware. The findings (Appendix B-6) indicated there was a significant decrease in the SSI rate from 1.1% in the pre-intervention group (n= 930) to 0.2% in the intervention group (n= 962) (p= 0.020). In the pre-intervention group (n= 930), of the positive SSIs, 10 were MSSA and 8 were MRSA. Of the intervention group (n= 962), two of the SSIs were MSSA and none were MRSA. The annual infection rate for the first year of the study in the pre-intervention group was 1.5% (n= 476) and 0.7% for the second year (n= 454). The annual infection rate for the first year of the study in the intervention group was 0.2% (n= 484) and 0.2% in the second year (n= 478).

This research study was valuable to a patient population that was not represented in the current literature. This population could not receive intranasal mupirocin before surgery because mupirocin needs to be applied for five days. Since this was emergent surgery, meeting this criteria was not possible. Therefore, intranasal povidone-iodine was a promising option as it only needs to be administered once before surgery. The results of this study demonstrated that intranasal povidone-iodine was not only effective in reducing the rate of SSIs, but also more cost effective. The researchers concluded that

adding the intranasal povidone-iodine to their decontamination protocol met statistical significance ($p= 0.020$), was cost effective, simple to use, and is widely available to medical professionals.

Cross Study Analysis

A cross study analysis was conducted and will be summarized next. Tables illustrating the findings and themes of each research study can be found in Appendix C.

A main outcome examined in this integrative review was whether the authors saw a reduction in SSIs following their intervention. Three of the six research studies explored the reduction of SSIs as one of their outcomes (Bebko et al., 2015; Phillips et al., 2014; Urias et al., 2018). All three authors found that their intervention, which consisted of daily CHG baths and intranasal povidone-iodine pre-operatively, resulted in a reduction of post-operative SSIs. Of these three authors, only one, Phillips et al. (2014), examined the use of CHG baths and intranasal povidone-iodine or CHG baths and intranasal mupirocin and impact on post-operative SSIs. These authors found that CHG baths and intranasal povidone-iodine was more effective at reducing SSIs than CHG baths and intranasal mupirocin.

The next outcome examined was the effectiveness of an intranasal decolonization protocol on the reduction of MRSA/MSSA. Research has shown that colonization with MRSA/MSSA in the nares is a risk factor for the development of a post-operative SSI. Therefore, the decolonization of the nares prior to surgery results in lower post-operative SSI rates. Three of the six studies (Anderson et al., 2015; Peng et al., 2018; Sai et al., 2015) studied the effects of a decontamination protocol on the reduction of nasal MRSA

and/or MSSA and all showed a reduction in nasal MRSA and/or MSSA. Anderson et al. (2015) analyzed the effects of intranasal povidone-iodine on MSSA and found that *S. aureus* growth was greatly reduced versus the control with the use of intranasal povidone-iodine. Peng et al. (2018) studied the effectiveness of intranasal povidone-iodine and CHG baths on the reduction of MSSA/MRSA and found a significant reduction of MSSA/MRSA post intervention. Sai et al. (2015) examined intranasal mupirocin with CHG baths versus intranasal povidone-iodine and CHG baths and their effectiveness against MRSA finding that the povidone-iodine group was not as successful as the mupirocin group, but it was effective in reducing the infection rate.

The final outcome examined was the cost effectiveness of intranasal mupirocin versus intranasal povidone-iodine. Four of the six research studies evaluated the cost of intranasal povidone-iodine versus intranasal mupirocin. All four studies concluded that the use of intranasal povidone iodine is much more cost effective than intranasal mupirocin (Bebko et al., 2015; Peng et al., 2018; Phillips et al., 2014; Urias et al., 2018). All authors noted the cost effectiveness to be due to the lower cost of povidone-iodine swabs and needing less povidone-iodine swabs versus mupirocin swabs.

There were several limitations noted in each study. All studies noted generalizability of the study as a limitation (Anderson et al., 2015; Bebko et al., 2015; Peng et al., 2018; Phillips et al., 2014; Sai et al., 2015; Urias et al., 2018) as they were conducted at one institution and with specific patient populations. One study noted that their positive results could be underestimated due to a short follow-up period (Bebko et al., 2015). A study by Peng et al. (2018) noted that they opted for culture swabs instead of PCR swabs due to cost, but the PCR swabs would have been more sensitive. Urias et

al. (2018) noted that they chose not to culture patients pre or post intervention to test the efficacy of the intranasal povidone-iodine. Five out of the six studies noted the need for more randomized control trials (Anderson et al., 2015; Bebko et al., 2015; Peng et al., 2018; Sai et al., 2015; Urias et al., 2018) as they were either retrospective, cross-sectional, or prospective. Finally, all authors noted the need for larger studies (Anderson et al., 2015; Bebko et al., 2015; Peng et al., 2018; Phillips et al., 2014; Sai et al., 2015; Urias et al., 2018).

Next, the summary and conclusions will be presented.

Summary and Conclusions

Surgical site infections have long been an unwanted complication of all types of surgical procedures. In the forefront of SSI research, many efforts have been focused on orthopedic surgeries. Mupirocin administered intranasally for five days before surgery has been a commonly used preventative to post-operative SSIs and has been shown to be effective by multitudes of clinical studies. However, there are concerns of rising mupirocin resistance, increased cost, and user error surrounding the use of intranasal mupirocin. For this reason, intranasal povidone-iodine is being investigated as another alternative. At this time, it does not have resistance, is cost effective, and is administered by a healthcare practitioner, significantly reducing user error.

The purpose of this integrative review was to evaluate the effectiveness of intranasal mupirocin versus intranasal povidone-iodine and its effectiveness in reducing nasal MRSA/MSSA colonization to prevent post-operative SSIs. After a thorough review of all relevant literature, research studies were compared to evaluate the effectiveness of intranasal povidone-iodine versus intranasal mupirocin on the rate of post-operative SSIs. By performing this comprehensive literature review, important research was compiled, and relevant outcomes were evaluated. The Iowa Model of Evidence-Based Practice was used to guide this integrative review. A thorough literature review was implemented which explored the use of intranasal povidone-iodine as an alternative to intranasal mupirocin in the prevention of SSIs. Following the literature review, an in-depth search on the use of intranasal mupirocin and intranasal povidone-iodine was explored. Search strategies and key words searched are located in the methods section. The PRISMA flow diagram was used to guide the selection of the research studies included in this

integrative review. All research studies that were used in this integrative review are examined in the results section. A brief review of those studies' findings will be presented next.

Phillips et al. (2014) published the only randomized control trial on the use of nasal mupirocin ointment versus intranasal povidone-iodine solution to date. The researchers found that the use of intranasal mupirocin was similar in efficacy to pre-operative intranasal povidone-iodine in preventing *S. aureus* SSI after arthroplasty and spine fusion surgery. Participants in the povidone-iodine intervention group were noted to have lower rates of treatment related symptoms and greater compliance as it was applied by a healthcare provider. Phillips et al. (2014) concluded that nasal povidone-iodine may be considered as an alternative to mupirocin to reduce the incidence of SSIs.

Anderson et al. (2015) performed a study to examine the efficacy of povidone-iodine in the prevention of MRSA infections in an ex vivo model, ex vivo human skin model, and in the anterior nares of human subjects. In all models studied, the povidone-iodine groups had significantly persistent reduced or undetectable bacterial densities, much greater than the mupirocin groups. The effectiveness of povidone-iodine on normal flora of the anterior nares in human subjects (n=70) was observed and it was discovered that *S. aureus* was significantly reduced from baseline level than the saline control subjects. It was concluded that the benefits of povidone-iodine should be considered in the prevention of SSIs as well as evolving mupirocin resistance and issues with medication compliance.

Bebko et al. (2015) performed a prospective clinical study on the effectiveness of a decontamination protocol consisting of chlorhexidine washcloths and chlorhexidine oral rinse the night before and the morning of orthopedic surgery in addition to intranasal povidone-iodine once the morning of surgery (n= 344). The control group consisted of patients who did not receive a decontamination protocol (n= 365). The results indicated that the SSI rate in the intervention group was significantly lower than that in the control group. Methicillin resistant staphylococcus aureus was identified as an independent predictor of developing an SSI. Povidone-iodine was supported as an attractive alternative to SSI prevention due to its efficacy, cost-effectiveness, and fewer concerns about antibiotic resistance.

Sai et al. (2015) performed a retrospective cohort study to evaluate the effectiveness of two decontamination protocols on positive MRSA carriers. The chlorhexidine group had a 56% success rate and the povidone-iodine group had a 23% success rate. The authors concluded that intranasal mupirocin was more effective than intranasal povidone-iodine, but they did note that the povidone-iodine was implemented on the complicated cases. While the researchers did not demonstrate a high success rate, they concluded that the effectiveness of decolonization on infection rates justified the continuation of a decontamination protocol.

Peng et al. (2018) conducted a prospective cross-sectional study to assess the prevalence of MSSA/MRSA in patients undergoing elective orthopedic surgery as well as to determine whether their current treatment protocols resulted in the successful decolonization of MSSA/MRSA. The results showed successful eradication of MRSA and significantly reduced colonization of MSSA. The researchers concluded that the use

of intranasal povidone-iodine was as effective as intranasal mupirocin and more cost effective.

Urias et al. (2018) performed a retrospective review evaluating the effectiveness of two decontamination protocols to reduce SSIs in urgent lower extremity repairs. The results were statistically significant, with a decrease in SSIs from 1.1% in the pre-intervention group to 0.2% in the intervention group, which included the use of povidone-iodine. The researchers recommended the use of intranasal povidone-iodine for its efficacy, cost-effectiveness, ease of use, and wide availability.

Throughout this integrative review, three out of the six studies showed increased effectiveness of povidone-iodine versus mupirocin in the reduction of SSIs (Bebko et al., 2015; Phillips et al. 2014; Urias et al., 2018). Three out of the six studies found a significant reduction in MRSA/MSSA colonization with the use of intranasal povidone-iodine versus mupirocin (Anderson et al., 2015; Peng et al., 2018; Urias et al. 2018). Overall, five out of the six studies found that statistically significant results proving that intranasal povidone-iodine was as effective or more effective than intranasal mupirocin (Anderson et al., 2015; Bebko et al., 2015; Peng et al., 2018; Phillips et al., 2014; Urias et al., 2018). One research study found a low success rate in the reduction of MRSA/MSSA with the use of intranasal povidone iodine, but did show effectiveness (Sai et al., 2015). Four of the six studies showed that intranasal povidone-iodine was more cost effective than intranasal mupirocin (Bebko et al., 2015; Peng et al., 2018, Phillips et al., 2014; Urias et al., 2018).

This review yielded largely positive evidence that the use of intranasal povidone iodine is effective in reducing the rate of SSIs, MRSA and MSSA. There are several limitations to this integrative review to be acknowledged. Unfortunately, there was only one randomized control trial found during the literature search, with the other studies being cross-sectional, prospective, or retrospective. Therefore, there is a need for larger randomized control trials to further support the current clinical evidence that intranasal povidone-iodine is just as effective, if not more effective, than intranasal mupirocin in the prevention of SSIs. Additionally, a majority of these research studies included chlorhexidine baths in their decolonization protocol, which may have attributed to the decrease in SSIs.

In conclusion, intranasal povidone-iodine is an attractive alternative to intranasal mupirocin in the fight against SSIs, as has been supported in the aforementioned research studies. The use of povidone-iodine has been shown to reduce SSIs, reduce and/or eradicate MRSA and MSSA, be more cost-effective, reduce mupirocin resistance, reduce non-compliance rates, and be user friendly. Intranasal povidone-iodine is a promising alternative to SSI prevention and consideration by medical professionals should be taken to implement this effective alternative.

Next, the recommendations and implications for advanced nursing practice will be presented.

Recommendations and Implications for Advanced Nursing Practice

At the forefront of healthcare decisions and implementations is evidence-based practice (EBP). Evidence based practice is the culmination of research studies that aim to demonstrate the most effective and safe treatment for patients and it is ever changing. One thing that a healthcare provider can always count on is always learning new information and improving upon current practice. What was new, top of the line research just a decade later will be outdated and modified per guidelines from new research. It is for this reason that advanced practice registered nurses (APRNs) must continue to evaluate the newest literature to provide the best care and most up to date treatment guidelines for their patients. This integrative review provided valuable, up to date information based on evidence and provides important guidelines for APRNs.

No matter which specialty an APRN practices in, preventing infection will be one of their top priorities. In the case of surgical patients, it is of utmost importance that the APRN take all possible measures to prevent a post-operative SSI from occurring. Current practice in the prevention of SSIs is up to the discretion of each individual provider and healthcare system. As noted in this integrative review, the use of intranasal povidone-iodine has proven to be just as, if not more, effective than intranasal mupirocin. As an APRN, their job is to give their patients the best care possible. This includes providing them with treatment regimens that are cost-effective, easy to use, effective, and are easy to comply with.

Advanced practice NPs are trained at the masters' level to be able to practice autonomously, which includes diagnosing, ordering labs and tests, and prescribing medications. They are also trained in the leadership role and in the ability to promote

change within institutions. This leadership role can be used to provide research, education, policy change, and implementation for healthcare workers. This is an important quality of the APRN which could be used to promote a policy change in an institution and implement the use of intranasal povidone-iodine administered pre-operatively to further reduce SSI rates.

In order to facilitate policy change, key interdisciplinary team members must be involved. Related to instituting intranasal povidone-iodine in the prevention of SSIs, key interdisciplinary members would include, but not limited to, nursing supervisors, surgeons, infection control, pharmacists, and nursing staff. It is the role of the APRN to present research and evidence-based findings and to provide education to these specialties to promote policy change. Once key team members have agreed to implementation of a new policy, the APRN has the ability develop the new policy and provide the education to implement the new policy change. This role is invaluable to promoting change and providing the best care for patients according to the most up to date research studies. The research compiled in this integrative review can be utilized to create a new policy in the fight against post-operative SSIs.

As previously noted in this integrative review, intranasal mupirocin requires application for five days before surgery which makes it harder for patients to comply with the treatment plan and furthermore, is not possible in more emergent surgeries. The use of intranasal povidone-iodine is an attractive alternative because it only needs to be applied once before surgery, therefore it's able to be administered on an urgent basis, and it is applied by a nurse before surgery therefore, patient compliance is eliminated as a risk

factor. The APRN will be able to provide research and education to nurses on the proper administration of intranasal povidone-iodine.

Education is a valuable role of the APRN. Of specific value are the years of experience the NP has in also being a registered nurse. Due to this background, the NP has the valuable trait of being more like a peer to the nurse rather than an authority figure. This puts the NP in a strategic position to be able to relate to the nurse and gain valuable trust, enabling an open relationship where the nurse and APRN can feel comfortable and free to learn from each other. This trust and understanding, as well as the advanced education of the APRN, provides a conducive environment for education. With the APRN's advanced degree, they are also able to educate other healthcare providers. This includes fellow mid-levels, physicians, and surgeons.

Before education is administered, research must occur. Research is the start of all evidence-based practice and is the avenue for policy change. Another valuable asset of the APRN is being able to attain, interpret, and disseminate valuable new information attained from the latest research studies. Evidence-based practice is constantly evolving and will always remain an important part of the APRN role. As noted in this integrative review, there needs to be further research on the use of intranasal povidone-iodine versus intranasal mupirocin. Once the APRN gathers and critiques all of the evidence from research, it can be disseminated to colleagues and then assist in providing education and training for practice.

While this integrative review showed very promising results in using intranasal povidone-iodine to prevent SSIs, further research is needed. Unfortunately, there is only

one randomized control trial evaluating the effectiveness of intranasal povidone-iodine versus intranasal mupirocin in the prevention of SSIs. Therefore, further randomized control trials and studies with larger sample sizes are needed. The APRN is in a valuable position to be able to implement a research study at their own institution and create definitive practice guidelines.

An important consideration for APRNs is the cost of the medications they order for their patients. Patients have many different economic backgrounds, and some don't even have insurance. In order for patients to take their medications, they need to first be able to afford them. Several articles in this integrative review found that intranasal povidone-iodine is more cost-effective than intranasal mupirocin, which is an important consideration to make when ordering medications to prevent SSIs.

When the NP is choosing a better alternative for patients in the prevention of SSIs, it is important that the treatment they choose is effective. Of the six articles presented in this integrative review, all showed the effectiveness of intranasal povidone-iodine. Despite the results presented in the article by Sai et al. (2015) that showed a low success rate at the eradication of MSSA/MRSA decolonization, intranasal povidone-iodine was effective in lowering infection rates. With all of these research studies included in this review showing positive results, it is recommended that the use of intranasal povidone-iodine be considered by the APRN for use in the prevention of SSIs.

Advanced practice registered nurses are in the unique position of having the background of a registered nurse with the capabilities of prescribing and improving patient outcomes. This puts them in an advantageous position to advocate for change and

have easy access to educating nurses and medical professionals. The APRN is a valuable resource for educating nurses and providers on current evidence-based practice and implementing new practice guidelines and policies for their patient populations.

References

- Alexander, J. W., Solomkin, J. S., & Edwards, M. J. (2011). Updated recommendations for control of surgical site infections. *Annals of Surgery*, 263(6), 1082-1093. doi: 10.1097/SLA.0b013e31821175f8
- Alfonso-Sanchez, J. L., Martinez, I. M., Martin-Moreno, J. M., Gonzalez, R. S., & Botia, F. (2017). Analyzing the risk factors influencing surgical site infections: the site of environmental factors. *Can J Surg*, 60(3), 155-161. doi: 10.1503/cjs.017916
- Al-Tamimi, M., Himsawi, N., Abu-Raideh, J., Jazar, D. A., Al-Jawaldeh, H., Al Haj Mahmoud, S., Hijjawi, N., & Hawamdeh, H. (2018). Nasal colonization by methicillin-sensitive and methicillin-resistant staphylococcus aureus among medical students. *The Journal of Infection in Developing Countries*, 12(5), 326-335. doi: 10.3855/jidc.9908
- Anderson, D. J., Podgorny, K., Berrios-Torres, S. I., Bratzler, D. W., Dellinger, E. P., Greene, L., Nyquist, A. C., Saiman, L., Yokoe, D. S., Maragakis, L. L., & Kaye K. S. (2014). Strategies to prevent surgical site infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol*, 36(6), 605-627. doi: 10.1086/676022
- Anderson, M. J., David, M. L., Scholz, M., Bull, S. J., Morse, D., Hulse-Stevens, M., & Peterson, M. L. (2015). Efficacy of skin and nasal povidone-iodine preparation against mupirocin-resistant staphylococcus aureus and s. aureus within the anterior nares. *Antimicrobial Agents and Chemotherapy*, 59(5), 2765-2773.

- Andersson, A. E., Bergh, I., Karlsson, J., Eriksson, B. I., & Nilsson, K. (2012). The application of evidence-based measures to reduce surgical site infections during orthopedic surgery-report of a single-center experience in Sweden. *Patient Safety in Surgery*, 6(11), 1-8.
- Bandaru, N. R., Rao, R. A., Prasad, V. K., & Murty, R. (2012). A prospective study of postoperative wound infections in a teaching hospital of rural setup. *Journal of Clinical & Diagnostic Research*, 6(7), 1266-1271.
- Bebko, S. P., Green, D. M., & Awad, S. S. (2015). Effect of a preoperative decontamination protocol on surgical site infections in patients undergoing elective orthopedic surgery with hardware implantation. *JAMA Surgery*, 150(5), 390-395. doi: 10.1001/jamasurg.2014.3480
- Berrios-Torres, S. I., Umscheid, C. A. & Bratzler, D. W. (2017). Centers for disease control and prevention guideline for the prevention of surgical site infection, 2017. *JAMA Surg*, 152(8), 784-791. doi: 10.1001/jamasurg.2017.0904
- Bonnel, W. & Smith, K. V. (2018). *Proposal writing for clinical nursing and DNP projects* (2nd ed.). New York, NY: Springer Publishing Company.
- Brown, C. G. (2014). The Iowa Model of Evidence-Based Practice to promote quality care: An illustrated example in oncology nursing. *Clinical Journal of Oncology Nursing*, 18(2), 157-159. doi: 10.1188/14.CJON.157-159
- Burks, I. R. (1998). Povidone-iodine solution in wound treatment. *Physical Therapy*, 78(2), 212-218.

Campbell, N. & Campbell, D. (2013). Evaluation of a non-adherent, povidone-iodine dressing in a case series of chronic wounds. *Journal of Wound Care*, 22(8), 401-406.

Centers for Disease Control and Prevention (CDC) (2019). *National Center for Emerging and Zoonotic Infectious Diseases Division of Healthcare Quality Promotion*.

Retrieved from: www.cdc.gov

Cheng, K., Li, J., Kong, Q., Wang, C., Ye, N., & Xia, G. (2015). Risk factors for surgical site infection in a teaching hospital: A prospective study of 1, 138 patients. *Patient Preference and Adherence*, 9, 1171-1177.

Hogan, B., Rakotozandrindrainy, R., Al-Emran, H., Dekker, D., Hahn, A., Jaeger, A., Poppert, S., Frickmann, H., Matthias Hagen, R., Micheel, V., Crusius, S., Heriniaina, J. N., Rakotondrainiarivelo, J. P., Razafindrabe, T., May, J., & Schwarz, N. G. (2016). Prevalence of nasal colonization by methicillin-sensitive and methicillin-resistant staphylococcus aureus among healthcare workers and students in madagascar. *BMC Infectious diseases*, 16(420), 1-9. doi: 10.1186/s12879-016-1733-6

Hussain, M. S., Naqvi, A., & Sharaz, M. (2019). Methicillin resistant staphylococcus aureus (MRSA); Prevalence and susceptibility pattern of (MRSA) isolated from pus in tertiary care of district hospital of Rahim Yar Khan. *The Professional Medical Journal*, 26(1), 122-127. doi: 10.29309/TPMJ/2019.26.01.2510

Iowa Model Collaborative (2017). Iowa Model of Evidence-Based Practice: Revisions and validation. *Worldviews on Evidence-Based nursing*, 14(3), 175-182.

- Kavanagh, K. T., Abusalem, S., & Calderon, L. E. (2017). The incidence of MRSA infections in the United States: Is a more comprehensive tracking system needed? *Antimicrobial Resistance and Infection Control*, 6(34), 1-6. doi: 10.1186/s13756-017-0193-0
- Kim, M. W., Greenfield, B. K., Snyder, R. E., Steinmaus, C. M., & Riley, L. W. (2018). The association between community-associated staphylococcus aureus colonization and disease: A meta-analysis. *BMC Infectious Diseases*, 18(86), 1-11. doi: <https://doi.org/10.1186/s12879-018-2990-3>
- Ko, W., Lazenby, W. D., Zelano, J. A., Isom, O. W., & Krieger, K. H. (1992). Effects of shaving methods and intraoperative irrigation on suppurative mediastinitis after bypass operations. *Ann Thoracic Surg*, 53(2), 301-305. doi: 10.1016/0003-4975(92)91337-9
- Kourtis, A. P., Hatfield, K., Baggs, J., Mu, Y., See, I., Epton, E., Nadle, J., Kainer, M. A., Dumyati, G., Petit, S., & Ray, S. M. (2019). *Morbidity and Mortality Weekly Report*, 68(9), 214-219.
- Loftus, R. W., Dexter, F., & Robinson, A. D. M. (2018). High-risk staphylococcus aureus transmission in the operating room: A call for widespread improvements in perioperative hand hygiene and patient decolonization practices. *American Journal of Infection Control*, 46(10), 1134-1141. doi: <https://doi.org/10.1016/j.ajic.2018.04.211>
- Loftus, R. W., Dexter, F., & Robinson, A. D. M. (2018). Methicillin-resistant staphylococcus aureus has greater risk of transmission in the operating room than

methicillin-sensitive *S aureus*. *American Journal of Infection Control*, 46, 520-525. doi: <https://doi.org/10.1016/j.ajic.2017.11.002>

Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & The PRISMA Group. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLOS Medicine*, 6(7), 1-6.

National Hospital Inpatient Quality Measures Specifications Manual (NHIQMSM). (2014). Retrieved from: www.qualitynet.org

Owens, C. D. & Stoessel, K. (2008). Surgical site infections: epidemiology, microbiology, and prevention. *Journal of Hospital Infection*, 70(52), 3-10.

Peng, H. M., Wang, L. C., Zhai, J. L., Weng, X. S., Feng, B., & Wang, W. (2018). Effectiveness of preoperative decolonization with nasal povidone iodine in Chinese patients undergoing elective orthopedic surgery: A prospective cross-sectional study. *Brazilian Journal of Medical and Biological Research*, 51(2), 1-6. doi: 10.1590/1414-431X20176736

Perl, T. M., Cullen, J. J., Wenzel, R. P., Zimmerman, M. B., Pfaller, M. A., Sheppard, D., Twombly, J., French, P. P., & Herwaldt, L. A. (2002). Intranasal mupirocin to prevent postoperative staphylococcus aureus infections. *The New England Journal of Medicine*, 346, 1871-1877. doi: 10.1056/NEJMoa003069

Phillips, M., Rosenberg, A., Shopsis, B., Cuff, G., Skeete, F., Foti, A., Kraemer, K., Inglis, K., Press, B., & Bosco, J. (2014). Preventing surgical site infections: A randomized, open-label trial of nasal mupirocin ointment and nasal povidone

iodine solution. *Infection Control Hospital of Epidemiology*, 35(7), 826-832. doi: 10.1086/676872

- Piechota, M., Kot, B., Frankowska-Maciejewska, A., Gruzewska, A., & Wozniak-Kosek, A. (2018). Biofilm formation by methicillin-resistant and methicillin-sensitive staphylococcus aureus strains from hospitalized patients in poland. *BioMed Research International*, 2018, 1-7. doi: <https://doi.org/10.1155/2018/4657396>
- Polit, D. F. & Beck, C. T. (2017). *Nursing research: generating and assessing evidence for nursing practice*. (10th ed.). Philadelphia, PA: Wolters Kluwer.
- Ramos, N., Skeete, F., Haas, J. P., Hutzler, L., Slover, J., & Phillips, M. (2011). Surgical site infection prevention initiative patient attitude and compliance. *Bulletin of the NYU Hospital for Joint Diseases*, 69(4), 312-318.
- Rosenberger, L. H., Politano, A. D., & Sawyer, R. G. (2011). The surgical care improvement project and prevention of post-operative infection, including surgical site infection. *Surgical Infections*, 12(3), 163-168. doi: 10.1089/sur.2010.083
- Sai, N., Laurent, C., Strale, H., Denis, O., & Byl, B. (2015). Efficacy of the decolonization of methicillin-resistant staphylococcus aureus carriers in clinical practice. *Antimicrobial Resistance and Infection Control*, 4(56), 1-8. doi: 10.1186/s/13756-015-0096-x
- Sakr, J., Bregeon, F., Mege, J. L., Rolain, J. M., & Blin, O. (2018). Staphylococcus aureus nasal colonization: An update on mechanisms, epidemiology, risk factors,

and subsequent infections. *Frontiers in Microbiology*, 9(2419), 1-15. doi:
10.3389/fmicb.2018.02419

Septimus, E. J. & Schweizer, M. L. (2016). Decolonization in prevention of health care-associated infections. *American Society for Microbiology*, 29(2), 201-222.

Sfeir, M., Obeid, Y., Eid, C., Saliby, M., Farra, A., Farhat, H., & Mokhbat, J. E. (2014). Prevalence of staphylococcus aureus methicillin-sensitive and methicillin-resistant nasal and pharyngeal colonization in outpatients in Lebanon. *American Journal of Infection Control*, 42, 160-163. doi:
<http://dx.doi.org/10.1016/j.ajic.2013.08.008>

Stone, P. W., Kunches, L., & Hirschhorn, L. (2009). Cost of hospital-associated infections in massachusetts. *American Journal of Infection Control*, 37(3), 210-214. doi:
<https://doi.org/10.1016/j.ajic.2008.07.011>

Surahio, A. R., Talpur, A. A., Memon, A. S., Junejo, A., & Laghari, A. A. (2017). Surgical site infections: Culture & sensitivity pattern of micro-organisms isolated in a tertiary care hospital. *The Professional Medical Journal*, 24(1), 57-63.

Uchiyama, S., Dahesh, S., Nizet, V., & Kessler, J. (2019). Enhanced topical delivery of non-complexed molecular iodine for methicillin-resistant staphylococcus aureus decolonization. *International Journal of Pharmaceutics*, 554, 81-86. doi:
<https://doi.org/10.1016/j.ijpharm.2018.11.004>

Urias, D. S., Varghese, M., Simunich, T., Morrissey, S., & Dumire, R. (2018).

Preoperative decolonization to reduce infections in urgent lower extremity repairs.

European Journal of Trauma and Emergency Surgery, 44, 787-793.

Waryah, C. B., Gogoi-Tiwari, J., Wells, K., Eto, K. Y., Masoumi, E., Costantino, P.,

Kotiw, M., & Mukkur, T. (2016). Diversity of virulence factors associated with

west Australian methicillin-sensitive staphylococcus aureus isolates of human

origin. *BioMed Research International*, 2016, 1-10. doi:

<http://dx.doi.org/10.1155/2016/8651918>

Yi, J., Liang, H., Song, R., Xia, H., & Huang, Y. (2018). Maintaining intraoperative

normothermia reduces blood loss in patients undergoing major operations: A pilot

randomized controlled clinical trial. *BMC Anesthesiology*, 18(126), 1-7. doi:

<https://doi.org/10.1186/s1287-018-0582-9>

Zinn, J., Jenkins, J. B., Swofford, V., Harrelson, B., & McCarter, S. (2010).

Intraoperative patient skin prep agents: Is there a difference? *AORN Journal*,

92(6), 662-671. doi: 10.1016/j.aorn.2010.07.016

Appendix A-1

Phillips, M., Rosenberg, A., Shopsin, B., Cuff, G., Skeete, F., Foti, A., Kraemer, K., Inglima, K., Press, B., & Bosco, J. (2014). Preventing surgical site infections: A randomized, open-label trial of nasal mupirocin ointment and nasal povidone iodine solution. *Infection Control Hosp Epidemiol*, 35(7), 826-832

Title	<ul style="list-style-type: none"> Is the title a good one, succinctly suggesting key variables and the study population? 	The title was good in regard to key variables but only made reference to the study population broadly, stating that it was a surgical population.
Abstract	<ul style="list-style-type: none"> Did the abstract clearly and concisely summarize the main features of the report (problem, methods, results, conclusions)? 	The abstract was well written and clearly and concisely summarized the main features of the report. The background clearly stated the problem and the methods; results, and conclusions sections were very informative and included key information in the study.
Introduction Statement of the Problem	<ul style="list-style-type: none"> Was the problem stated unambiguously, and was it easy to identify? Is the problem statement build a persuasive argument for the new study? Was there a good match between the research problem and the methods used –that is, was a 	The introduction section introduced the problem unambiguously and it was easy to identify in the opening paragraph with a statement about surgical site infection rates. A persuasive argument was built for this study by providing statistics and current

	quantitative approach appropriate?	practices that the mentioned hospital performs. There was a good match between the research problem and the methods used. A quantitative approach was most certainly appropriate in this study because they were collecting data on infection rates, cost and compliance.
Hypotheses or Research Questions	<ul style="list-style-type: none"> • Were research questions and/or hypotheses explicitly stated? If not, was their absence justified? • Were questions and hypotheses appropriately worded, with clear specification of key variables and the study population? • Were the questions/hypotheses consistent with existing knowledge? 	A research question was not explicitly stated but a hypothesis was explicitly stated at the end of the introduction section. The hypothesis was expertly and appropriately worded with very clear and concise specification of key variables. Once again, the study population was only broadly mentioned as surgical patients, but not specifically which type of surgical patients. The hypothesis was consistent with existing knowledge.
Literature Review	<ul style="list-style-type: none"> • Was the literature review up-to-date and based mainly on primary sources? • Did the review provide a state-of- 	Not all of the literature was up to date in the literature review. Many of the early cited articles were from 2011-

	<p>the-art synthesis of evidence on the problem?</p> <ul style="list-style-type: none"> • Did the literature review provide a strong basis for the new study? 	<p>2013. As the literature review went on it started to get specific about the use of mupirocin and the reduction of surgical site infections and some of these references dated back as far as 1995. All sources referenced were primary sources. The review was fairly brief and could have been improved. Despite this, the literature review certainly provided a strong basis for the new study.</p>
<p>Conceptual/Theoretical Framework</p>	<ul style="list-style-type: none"> • Were key concepts adequately defined conceptually? • Was a conceptual/theoretical framework articulated—and, if so, was it appropriate? If not, is the absence of a framework justified? • Were the questions/hypotheses consistent with the framework? 	<p>Key concepts were adequately defined conceptually in this study. There was no conceptual/theoretical framework utilized in this study. The absence of a framework was justified as this quantitative study had a nice flow and was neatly summarized in various tables.</p>
<p>Method</p> <p>Protection of Human Rights</p>	<ul style="list-style-type: none"> • Were appropriate procedures used to safeguard the rights of study participants? • Was the study externally reviewed by an IRB/ethics 	<p>Appropriate procedures were used to safeguard the rights of study participants. The study was reviewed by the institutional review board at</p>

	<p>review board?</p> <ul style="list-style-type: none"> • Was the study designed to minimize risks and maximize benefits to participants? 	<p>their institution and informed consent was obtained from all study participants. The study was designed to minimize risks and maximize benefits to participants. Patients received verbal and written instructions and had access to a 24/7 phone number in case they had study treatment related questions.</p>
Research Design	<ul style="list-style-type: none"> • Was the most rigorous design used, given the study purpose? • Were appropriate comparisons made to enhance interpretability of the findings? • Was the number of data collection points appropriate? • Did the design minimize biases and threats to the internal, construct, and external validity of the study (e.g., was blinding used, was attrition minimized)? 	<p>The research design was an investigator initiated, open label, randomized trial study. This was the most rigorous design that could have been used. Appropriate comparisons were made to enhance the interpretability of the findings. There were three data collection outcomes: infection rates; cost; and patient compliance, which were appropriate for this study. The design did minimize biases and threats to the internal construct and external validity of the study by being randomized to either the mupirocin</p>

		or povidone iodine treatment groups. Patients were not blinded to their treatment group as they were aware if they were receiving mupirocin or povidone iodine.
Population and Sample	<ul style="list-style-type: none"> • Was the population identified? Was the sample described in sufficient detail? • Was the best possible sampling design used to enhance the sample's representativeness? Were sampling biases minimized? • Was the sample size based on a power analysis? 	<p>The population was identified in terms of which type of surgery they were undergoing. There was also a chart on demographic characteristics describing the age, sex, race, and ethnicity of the study participants. The best possible sampling design was used to enhance the sample's representativeness. Sampling biases were minimized. In the statistical analysis section there was mention of the sample size being based on a power analysis; however, this study was unable to reach the goal sample size.</p>
Data Collection and Measurement	<ul style="list-style-type: none"> • Were the operational and conceptual definitions congruent? • Were key variables measured using an appropriate method (e.g., interviews, 	<p>The operational and conceptual definitions were congruent in this study. The key variables were measured appropriately using</p>

	<p>observations, and so on)?</p> <ul style="list-style-type: none"> • Were specific instruments adequately described and were they good choices, given the study population and the variables being studied? • Did the report provide evidence that the data collection methods yielded data that were reliable, valid and responsive? 	<p>statistical analysis for the quantitative study. Specific instruments were adequately described and were good choices given this study population. Categorical variables were analyzed using Fisher's exact test. The report provides some evidence that the data collection methods yielded data that was reliable, valid, and responsive by utilizing practitioners who were blinded to study participation and receipt of study treatment. Also, microbiology reports, hospital readmissions and reports from other healthcare facilities were used.</p>
Procedures	<ul style="list-style-type: none"> • If there was an intervention, was it adequately described, and was it rigorously developed and implemented? Did most participants allocated to the intervention group actually receive it? Was there evidence of intervention fidelity? 	<p>The intervention of using mupirocin versus povidone iodine was certainly adequately described and was rigorously developed and implemented. All participants allocated to the intervention group received either mupirocin or</p>

	<ul style="list-style-type: none"> • Were data collected in a manner that minimized bias? Were the staff who collected data appropriately trained? 	<p>povidone iodine. There was no evidence of intervention fidelity indicated. Data were collected in a manner that minimized bias by practitioners being blinded to the study. The staff appeared to have been adequately trained on data collection as they were from Infection Prevention and Control.</p>
Data Analysis	<ul style="list-style-type: none"> • Were analyses undertaken to address each research question or test each hypothesis? • Were appropriate statistical methods used, given the level of measurement of the variables, number of groups being compared, and assumptions of the texts? • Was a powerful analytic method used? (e.g., did the analysis help to control for confounding variables)? • Were type I and Type II errors avoided or minimized? • In intervention studies, was an intention-to-treat analysis performed? 	<p>There was no mention of whether analyses were undertaken to address each research question or test each hypothesis. Statistical analysis was conducted using SAS version 9.1. Categorical variables were analyzed using Fisher's exact test. These seem appropriate statistical methods for this study. It is not clear in this study if the analysis helped to control for confounding variables. Type I and type II errors were avoided. An intention-to-treat analysis was performed. Problems of missing</p>

	<ul style="list-style-type: none"> • Were problems of missing values evaluated and adequately addressed? 	values were evaluated and adequately addressed.
Findings	<ul style="list-style-type: none"> • Was information about statistical significance presented? Was information about effect size and precision of estimates (confidence intervals) presented? • Were the findings adequately summarized, with good use of tables and figures? • Were findings reported in a manner that facilitates a meta-analysis, and with sufficient information needed for EBP? 	There was some information about statistical significance presented but there were no confidence intervals presented. The findings were adequately summarized with the use of multiple tables and figures. The findings were reported in a manner that facilitated a meta-analysis with sufficient information needed for EBP.
Discussion Interpretation of the Findings	<ul style="list-style-type: none"> • Were all major findings interpreted and discussed within the context of prior research and/or the study's conceptual framework? • Were casual inferences, if any, justified? • Was the issue of clinical significance discussed? • Were interpretations well-founded and consistent with the study's limitations? • Did the report address the issue of 	All major findings were interpreted and discussed within the context of prior research. The issue of clinical significance was discussed including infection rates, cost, and ease of use. Interpretations were well-founded and consistent with the study's limitations. The report did address the issue of generalizability of the findings given this study was only

	the generalizability of the findings?	at one institution and the results may not be applicable to other locations with different patient characteristics.
Implications/Recommendations	<ul style="list-style-type: none"> Did the researchers discuss the implications of the study for clinical practice or further research—and were those implications reasonable and complete? 	The researchers briefly discussed the implications of the study for further clinical practice stating that povidone iodine may be considered as an alternative to mupirocin and a component of a multifaceted approach to reduce surgical site infections.
General Issues Presentation	<ul style="list-style-type: none"> Was the report well-written, organized, and sufficiently detailed for critical analysis? In intervention studies, was a CONSORT flowchart provided to show the flow of participants in the study? Was the report written in a manner that makes the findings accessible to practicing nurses? 	The report was well written, organized, and fairly sufficiently detailed, although there could have been some improvement in this area. There was a CONSORT flowchart present to show the flow of the participants in the study. This report was written in a manner that makes the findings accessible to practicing nurses.
Researcher Credibility	<ul style="list-style-type: none"> Do the researchers' clinical, substantive, or methodologic qualifications and experience enhance confidence in the findings and their 	The researchers' clinical qualifications and experience enhanced confidence in the findings and their

	interpretation?	interpretation.
Summary Assessment	<ul style="list-style-type: none"> • Despite any limitations, do the study findings appear to be valid—do you have confidence in the truth value of the results? • Does the study contribute any meaningful evidence that can be used in nursing practice or that is useful to the nursing discipline? 	<p>While this was a small study at only one institution, there is confidence in the truth value of the results. This study definitely contributed meaningful evidence that can be used in nursing practice and/or the nursing discipline. Especially since povidone iodine only has to be used right before surgery, it greatly increases compliance and reduces cost.</p>

Appendix A-2

Anderson, M. J., David, M. L., Scholz, M., Bull, S. J., Morse, D., Hulse-Stevens, M., & Peterson, M. L. (2015). Efficacy of skin and nasal povidone-iodine preparation against mupirocin-resistant methicillin-resistant staphylococcus aureus and *S. aureus* within the anterior nares. *Antimicrobial Agents and Chemotherapy*, 59(5), 2765-2773.

Title	<ul style="list-style-type: none"> Is the title a good one, succinctly suggesting key variables and the study population? 	The title was good in terms of identifying the key variables, but it made no mention of the study population.
Abstract	<ul style="list-style-type: none"> Did the abstract clearly and concisely summarize the main features of the report (problem, methods, results, conclusions)? 	The abstract was great as it clearly and concisely summarized the main features of the report.
Introduction Statement of the Problem	<ul style="list-style-type: none"> Was the problem stated unambiguously, and was it easy to identify? Is the problem statement build a persuasive argument for the new study? Was there a good match between the research problem and the methods used –that is, was a quantitative approach appropriate? 	The problem was stated unambiguously and was very easy to identify and built a persuasive argument for a new study. There was a good match between the research problem and the methods as a quantitative study was appropriate in this situation to collect numerical data.
Hypotheses or Research Questions	<ul style="list-style-type: none"> Were research questions and/or hypotheses explicitly stated? If not, was their absence justified? 	The hypotheses were explicitly stated at the end of the introduction section and were clearly worded with

	<ul style="list-style-type: none"> • Were questions and hypotheses appropriately worded, with clear specification of key variables and the study population? • Were the questions/hypotheses consistent with existing knowledge? 	specification to key variables and the study population. The hypotheses were backed up with existing knowledge throughout the introduction section.
Literature Review	<ul style="list-style-type: none"> • Was the literature review up-to-date and based mainly on primary sources? • Did the review provide a state-of-the-art synthesis of evidence on the problem? • Did the literature review provide a strong basis for the new study? 	While a majority of the literature review was from the past ten years, there were some that dated as far back as twenty years. They were all primary sources. The literature review provided a state-of-the-art synthesis of the evidence on the problem and certainly provided a strong basis for the new study.
Conceptual/Theoretical Framework	<ul style="list-style-type: none"> • Were key concepts adequately defined conceptually? • Was a conceptual/theoretical framework articulated—and, if so, was it appropriate? If not, is the absence of a framework justified? • Were the questions/hypotheses consistent with the framework? 	There was no conceptual/theoretical framework articulated and its' absence was not addressed but the study was well articulated and had a good flow without it.
Method Protection of Human Rights	<ul style="list-style-type: none"> • Were appropriate procedures used to 	There was very little information

	<p>safe-guard the rights of study participants?</p> <ul style="list-style-type: none"> • Was the study externally reviewed by an IRB/ethics review board? • Was the study designed to minimize risks and maximize benefits to participants? 	<p>regarding appropriate procedures used to safeguard the rights of study participants. It was noted in the section regarding human skin explants that IRB review was not needed but under the human subjects' section there was no mention of an IRB review. While there was no mention of the study being designed to minimize risks and maximize benefits to participants, this study did involve minimal risk to study participants.</p>
Research Design	<ul style="list-style-type: none"> • Was the most rigorous design used, given the study purpose? • Were appropriate comparisons made to enhance interpretability of the findings? • Was the number of data collection points appropriate? • Did the design minimize biases and threats to the internal, construct, and external validity of the study (e.g., was blinding used, was attrition minimized)? 	<p>While this was clearly a quantitative study, there was no mention of which type of study this was. The authors simply referred to it as an analysis. This was appropriate as this was a very structured and organized study. There were five data collection points which was appropriate as there were five different interventions in this study. Threats to the internal construct and external validity of the study were minimized. In terms</p>

		of the human subjects, there was no mention of whether the design minimized biases and threats to external constructs/external validity.
Population and Sample	<ul style="list-style-type: none"> • Was the population identified? Was the sample described in sufficient detail? • Was the best possible sampling design used to enhance the sample's representativeness? Were sampling biases minimized? • Was the sample size based on a power analysis? 	While there were explicit details regarding bacterial growth and inoculation, ex vivo porcine vaginal mucosa, procurement culture of human skin explants, application of test formulations or comparators, mucin wash, and bacterial enumeration, the population of human subjects was not identified. Due to the lack of information, it was hard to determine if the best possible sampling design was used. There was no mention of a power analysis being used.
Data Collection and Measurement	<ul style="list-style-type: none"> • Were the operational and conceptual definitions congruent? • Were key variables measured using an appropriate method (e.g., interviews, observations, and so on)? 	The operational and conceptual definitions were congruent. Key variables were measured appropriately using observations and data collection. Specific instruments were briefly

	<ul style="list-style-type: none"> • Were specific instruments adequately described and were they good choices, given the study population and the variables being studied? • Did the report provide evidence that the data collection methods yielded data that were reliable, valid and responsive? 	mentioned and seemed to be good choices given the study population. The report provided evidence that the data collection methods yielded data that were reliable, valid, and responsive.
Procedures	<ul style="list-style-type: none"> • If there was an intervention, was it adequately described, and was it rigorously developed and implemented? Did most participants allocated to the intervention group actually receive it? Was there evidence of intervention fidelity? • Were data collected in a manner that minimized bias? Were the staff who collected data appropriately trained? 	There were several different interventions in this study and they were all very rigorously developed and implemented. There was a very detailed description of all interventions. Data were collected in a manner that minimized bias and there was mention of staff being appropriately trained.
Data Analysis	<ul style="list-style-type: none"> • Were analyses undertaken to address each research question or test each hypothesis? • Were appropriate statistical methods used, given the 	Statistical analyses were undertaken to address each hypothesis. The appropriate statistical methods were used in this study and included analysis of variance

	<p>level of measurement of the variables, number of groups being compared, and assumptions of the texts?</p> <ul style="list-style-type: none"> • Was a powerful analytic method used? (e.g., did the analysis help to control for confounding variables)? • Were type I and Type II errors avoided or minimized? • In intervention studies, was an intention-to-treat analysis performed? • Were problems of missing values evaluated and adequately addressed? 	<p>followed by Bonferroni's posttest using the GraphPad PRISM software. The student's t test was used to evaluate significant differences at each time point. There was no mention of whether an intention-to-treat analysis was performed. There were no missing values noted.</p>
Findings	<ul style="list-style-type: none"> • Was information about statistical significance presented? Was information about effect size and precision of estimates (confidence intervals) presented? • Were the findings adequately summarized, with good use of tables and figures? • Were findings reported in a manner that 	<p>Statistical significance was presented but there was no information about effect size and precision of estimates presented. There were some tables presented which adequately summarized the information. The findings were reported in a manner that facilitated a meta-analysis and with sufficient information needed for EBP.</p>

	facilitates a meta-analysis, and with sufficient information needed for EBP?	
Discussion Interpretation of the Findings	<ul style="list-style-type: none"> • Were all major findings interpreted and discussed within the context of prior research and/or the study's conceptual framework? • Were casual inferences, if any, justified? • Was the issue of clinical significance discussed? • Were interpretations well-founded and consistent with the study's limitations? • Did the report address the issue of the generalizability of the findings? 	The authors did a great job of discussing the findings within context of prior research. Casual inferences were justified. The issue of clinical significance was certainly discussed. Interpretations were well-founded and consistent with the study's limitations. There was no mention of the issue of the generalizability of the findings.
Implications/Recommendations	<ul style="list-style-type: none"> • Did the researchers discuss the implications of the study for clinical practice or further research—and were those implications reasonable and complete? 	The researchers did discuss the implications of the study for clinical practice and suggested that the use of povidone-iodine should be considered as an alternative for reducing the bioburden of anterior nares prior to surgery.
General Issues Presentation	<ul style="list-style-type: none"> • Was the report well-written, organized, and sufficiently detailed 	The report was well-written, organized, and sufficiently detailed for critical

	<p>for critical analysis?</p> <ul style="list-style-type: none"> • In intervention studies, was a CONSORT flowchart provided to show the flow of participants in the study? • Was the report written in a manner that makes the findings accessible to practicing nurses? 	<p>analysis. There was no CONSORT flowchart provided. The report was certainly written in a manner that makes the findings accessible to practicing nurses.</p>
Researcher Credibility	<ul style="list-style-type: none"> • Do the researchers' clinical, substantive, or methodologic qualifications and experience enhance confidence in the findings and their interpretation? 	<p>The credentials of the researchers were not noted but they belonged to the department of experimental and clinical pharmacology and were part of the 3M infection prevention division. They had experience in this field.</p>
Summary Assessment	<ul style="list-style-type: none"> • Despite any limitations, do the study findings appear to be valid—do you have confidence in the truth value of the results? • Does the study contribute any meaningful evidence that can be used in nursing practice or that is useful to the nursing discipline? 	<p>While the human portion of this study had a small sample size, there were also explant models which were inoculated in a controlled environment; therefore, the findings appeared to be valid. This study certainly contributed meaningful evidence that can be used in nursing practice, further supporting that the use of</p>

		povidone-iodine may be an alternative for reducing the bioburden of anterior nares prior to surgery.
--	--	--

Appendix A-3

Bebko, S. P., Green, D. M., & Awad, S. S. (2015). Effect of a preoperative decontamination protocol on surgical site infections in patients undergoing elective orthopedic surgery with hardware implantation. *JAMA Surgery*, 150(5), 390-395.

Title	<ul style="list-style-type: none"> Is the title a good one, succinctly suggesting key variables and the study population? 	The title did a good job of succinctly suggesting key variables and the study population.
Abstract	<ul style="list-style-type: none"> Did the abstract clearly and concisely summarize the main features of the report (problem, methods, results, conclusions)? 	The abstract summarized the main features of the report. There were separate sections for importance, objective, design, setting, participants, interventions, outcomes, measures, results, and conclusions.
Introduction Statement of the Problem	<ul style="list-style-type: none"> Was the problem stated unambiguously, and was it easy to identify? Is the problem statement build a persuasive argument for the new study? Was there a good match between the research problem and the methods used –that is, was a quantitative approach appropriate? 	In the introduction, the problem was stated unambiguously, and it was easy to identify. The problem statement did build a persuasive argument for the new study, citing the high cost and prevalence of surgical site infections. There was a good match between the research problem and the methods used and a quantitative study was appropriate.
Hypotheses or Research Questions	<ul style="list-style-type: none"> Were research questions and/or hypotheses explicitly 	The research question was explicitly stated. The

	<p>stated? If not, was their absence justified?</p> <ul style="list-style-type: none"> • Were questions and hypotheses appropriately worded, with clear specification of key variables and the study population? • Were the questions/hypotheses consistent with existing knowledge? 	<p>research question was appropriately worded with clear specification of key variables and the study population. The research question was consistent with existing knowledge.</p>
Literature Review	<ul style="list-style-type: none"> • Was the literature review up-to-date and based mainly on primary sources? • Did the review provide a state-of-the-art synthesis of evidence on the problem? • Did the literature review provide a strong basis for the new study? 	<p>The literature review consisted of primary sources dating back to the early 2000s. The review provided a state-of-the-art synthesis of evidence on the problem. The literature review provided a strong basis for the new study.</p>
Conceptual/Theoretical Framework	<ul style="list-style-type: none"> • Were key concepts adequately defined conceptually? • Was a conceptual/theoretical framework articulated—and, if so, was it appropriate? If not, is the absence of a framework justified? • Were the questions/hypotheses consistent with the framework? 	<p>The key concepts were adequately defined conceptually. There was no theoretical conceptual framework articulated. Its' absence is not justified in the study. The research question was consistent with the framework.</p>
Method Protection of Human Rights	<ul style="list-style-type: none"> • Were appropriate procedures used to safe-guard the rights of study 	<p>Appropriate procedures were used to safeguard the rights of the study</p>

	<p>participants?</p> <ul style="list-style-type: none"> • Was the study externally reviewed by an IRB/ethics review board? • Was the study designed to minimize risks and maximize benefits to participants? 	<p>participants. The study was approved by the institutional review board at the Michael E. DeBakey Veterans Affairs Medical Center. As this was a retrospective review of de-identified data, oral or written informed consent was waived. This study was designed to minimize the risks and maximize the benefits to participants.</p>
Research Design	<ul style="list-style-type: none"> • Was the most rigorous design used, given the study purpose? • Were appropriate comparisons made to enhance interpretability of the findings? • Was the number of data collection points appropriate? • Did the design minimize biases and threats to the internal, construct, and external validity of the study (e.g., was blinding used, was attrition minimized)? 	<p>The study used a quantitative design and this was appropriate given the study purpose. The study was described as a prospective clinical study. Appropriate comparisons were made to enhance interpretability of the findings. The number of data collection points was appropriate. The design minimized biases and threats to the internal, construct, and external validity of the study.</p>
Population and Sample	<ul style="list-style-type: none"> • Was the population identified? Was the sample described in sufficient detail? • Was the best possible sampling 	<p>The population was identified as patients undergoing elective orthopedic surgery with hardware implantation</p>

	<p>design used to enhance the sample's representativeness? Were sampling biases minimized?</p> <ul style="list-style-type: none"> Was the sample size based on a power analysis? 	<p>between October 1, 2012 through December 31, 2013. The best possible sampling design was used to enhance the sample's representativeness and sampling biases were minimized. There was no mention of a power analysis being used to determine the sample size. All patients that met the criteria during the specified period of time were utilized.</p>
Data Collection and Measurement	<ul style="list-style-type: none"> Were the operational and conceptual definitions congruent? Were key variables measured using an appropriate method (e.g., interviews, observations, and so on)? Were specific instruments adequately described and were they good choices, given the study population and the variables being studied? Did the report provide evidence that the data collection methods yielded data that were reliable, valid and responsive? 	<p>The operational and conceptual definitions were congruent. Key variables were measured appropriately using descriptive statistics for data collection and statistical analysis. The instruments were adequately described and were good choices given the study population and the variables being studied. The report did provide evidence that the data collection methods yielded data that were reliable, valid, and responsive.</p>
Procedures	<ul style="list-style-type: none"> If there was an intervention, was it 	<p>There was an intervention and it</p>

	<p>adequately described, and was it rigorously developed and implemented? Did most participants allocated to the intervention group actually receive it? Was there evidence of intervention fidelity?</p> <ul style="list-style-type: none"> • Were data collected in a manner that minimized bias? Were the staff who collected data appropriately trained? 	<p>was adequately described, rigorously developed, and implemented. All participants allocated to the intervention group actually received it. There was evidence of intervention fidelity. The researchers were blinded to patient identifiers; therefore bias was minimized. The staff who collected the data were appropriately trained.</p>
Data Analysis	<ul style="list-style-type: none"> • Were analyses undertaken to address each research question or test each hypothesis? • Were appropriate statistical methods used, given the level of measurement of the variables, number of groups being compared, and assumptions of the texts? • Was a powerful analytic method used? (e.g., did the analysis help to control for confounding variables)? • Were type I and Type II errors avoided or minimized? • In intervention studies, was an 	<p>Analyses were undertaken to address each research question. The univariate analysis was performed using a two-sided t test, the Pearson X2 test, and the Fisher exact test and it was performed at two levels. These were appropriate for the level of measurement of the variables, number of groups being compared, and the assumptions of the texts. Type I and II errors were avoided. There was no mention of an intention-to-treat analysis being performed. Problems of missing values</p>

	<p>intention-to-treat analysis performed?</p> <ul style="list-style-type: none"> • Were problems of missing values evaluated and adequately addressed? 	<p>were evaluated and adequately addressed.</p>
Findings	<ul style="list-style-type: none"> • Was information about statistical significance presented? Was information about effect size and precision of estimates (confidence intervals) presented? • Were the findings adequately summarized, with good use of tables and figures? • Were findings reported in a manner that facilitates a meta-analysis, and with sufficient information needed for EBP? 	<p>Information about statistical significance was presented. Information about effect size and precision of estimates were presented. Findings were adequately summarized with great use of tables and figures. Findings were reported in a manner that facilitates a meta-analysis with sufficient information needed for EBP.</p>
Discussion Interpretation of the Findings	<ul style="list-style-type: none"> • Were all major findings interpreted and discussed within the context of prior research and/or the study's conceptual framework? • Were casual inferences, if any, justified? • Was the issue of clinical significance discussed? • Were interpretations well-founded and consistent with the study's limitations? 	<p>All major findings were interpreted and discussed within the context of prior research. Casual inferences were justified. The issue of clinical significance was discussed. The interpretations were well-founded and consistent with the study's limitations. The report did address the issue of the generalizability</p>

	<ul style="list-style-type: none"> Did the report address the issue of the generalizability of the findings? 	of the findings.
Implications/Recommendations	<ul style="list-style-type: none"> Did the researchers discuss the implications of the study for clinical practice or further research—and were those implications reasonable and complete? 	The researchers discussed the implications of the study for clinical practice or further research. They stated that further studies should be performed including large-scale randomized controlled clinical trials and additional studies to validate the application of the decontamination protocol to other services implanting hardware.
General Issues Presentation	<ul style="list-style-type: none"> Was the report well-written, organized, and sufficiently detailed for critical analysis? In intervention studies, was a CONSORT flowchart provided to show the flow of participants in the study? Was the report written in a manner that makes the findings accessible to practicing nurses? 	The report was well-written, organized, and sufficiently detailed for critical analysis. There was a CONSORT flowchart provided to show the flow of participants in the study. The report was certainly written in a manner that makes the findings accessible to practicing nurses.
Researcher Credibility	<ul style="list-style-type: none"> Do the researchers' clinical, substantive, or methodologic qualifications and experience enhance confidence in the findings and their 	The researchers' clinical, substantive, and methodologic qualifications and experience enhance confidence in the findings and their

	interpretation?	interpretation. They were both from the department of surgery at Baylor College of Medicine.
Summary Assessment	<ul style="list-style-type: none"> • Despite any limitations, do the study findings appear to be valid—do you have confidence in the truth value of the results? • Does the study contribute any meaningful evidence that can be used in nursing practice or that is useful to the nursing discipline? 	Despite limitations to the study, the findings appear to be valid and there can be confidence in the truth value of the results. This study definitely contributed meaningful evidence that can be used in nursing practice and that is useful for the nursing discipline.

Appendix A-4

Sai, N., Laurent, C., Strale, H., Denis, O., & Byl, B. (2015). Efficacy of the decolonization of methicillin-resistant staphylococcus aureus carriers in clinical practice. *Antimicrobial Resistance and Infection Control*, 4(56), 1-8.

Title	<ul style="list-style-type: none"> Is the title a good one, succinctly suggesting key variables and the study population? 	The title was appropriate and succinctly suggested key variables and the study population.
Abstract	<ul style="list-style-type: none"> Did the abstract clearly and concisely summarize the main features of the report (problem, methods, results, conclusions)? 	The abstract clearly and concisely summarized the main features of the report.
Introduction Statement of the Problem	<ul style="list-style-type: none"> Was the problem stated unambiguously, and was it easy to identify? Is the problem statement build a persuasive argument for the new study? Was there a good match between the research problem and the methods used – that is, was a quantitative approach appropriate? 	The problem was stated unambiguously and was easy to identify. The problem statement did build a persuasive argument for a new study, citing that MRSA is associated with a high risk of acquiring MRSA infection during hospital stays. There was a good match between the research problem and the methods use; a quantitative study was appropriate.
Hypotheses or Research Questions	<ul style="list-style-type: none"> Were research questions and/or hypotheses explicitly stated? If not, was their absence justified? Were questions and hypotheses 	The research questions were explicitly stated, appropriately worded, and with clear specification of key variables and the study population, which was identified as

	<p>appropriately worded, with clear specification of key variables and the study population?</p> <ul style="list-style-type: none"> • Were the questions/hypotheses consistent with existing knowledge? 	<p>hospitalized patients. The research questions were certainly consistent with existing knowledge.</p>
Literature Review	<ul style="list-style-type: none"> • Was the literature review up-to-date and based mainly on primary sources? • Did the review provide a state-of-the-art synthesis of evidence on the problem? • Did the literature review provide a strong basis for the new study? 	<p>The literature review is fairly up-to-date with most of the data being from the early to mid-2000s. The review did provide a state-of-the-art synthesis of the evidence on the problem. The literature review provided a strong basis for the new study citing there was little information about the rate of success of decolonization strategies under real-life conditions.</p>
Conceptual/Theoretical Framework	<ul style="list-style-type: none"> • Were key concepts adequately defined conceptually? • Was a conceptual/theoretical framework articulated—and, if so, was it appropriate? If not, is the absence of a framework justified? • Were the 	<p>Key concepts were adequately defined conceptually. There was no theoretical framework articulated in this study and its absence is justified in this quantitative study.</p>

	questions/hypotheses consistent with the framework?	
Method Protection of Human Rights	<ul style="list-style-type: none"> • Were appropriate procedures used to safe-guard the rights of study participants? • Was the study externally reviewed by an IRB/ethics review board? • Was the study designed to minimize risks and maximize benefits to participants? 	Appropriate procedures were used to safeguard the rights of study participants. There was no external review by an IRB/ethics review board mentioned. The study was designed to minimize risks and maximize benefits to participants.
Research Design	<ul style="list-style-type: none"> • Was the most rigorous design used, given the study purpose? • Were appropriate comparisons made to enhance interpretability of the findings? • Was the number of data collection points appropriate? • Did the design minimize biases and threats to the internal, construct, and external validity of the study (e.g., was blinding used, was attrition minimized)? 	Given the study purpose, the most rigorous design was used. Appropriate comparisons were made to enhance interpretability of the findings. The number of data collection points was appropriate for this study. The design certainly minimized biases and threats to internal, construct, and external validity of the study.
Population and Sample	<ul style="list-style-type: none"> • Was the population 	The population was identified as inpatients

	<p>identified? Was the sample described in sufficient detail?</p> <ul style="list-style-type: none"> • Was the best possible sampling design used to enhance the sample's representativeness? Were sampling biases minimized? • Was the sample size based on a power analysis? 	<p>in an 864-bed academic hospital in Belgium. The best possible sampling design was used to enhance the sample's representativeness. Sampling biases were minimized. There was no mention of a power analysis. It appears the sample size was based on a convenience sample. The study population consisted of all patients diagnosed as MRSA-positive between January 2006 and June 2010.</p>
Data Collection and Measurement	<ul style="list-style-type: none"> • Were the operational and conceptual definitions congruent? • Were key variables measured using an appropriate method (e.g., interviews, observations, and so on)? • Were specific instruments adequately described and were they good choices, given the study population and the variables being studied? • Did the report provide evidence that the data collection methods yielded 	<p>The operational and conceptual definitions were congruent. Key variables were measured using a data collection tool and statistical analysis, appropriate for this quantitative study. The authors created their own data collection tool, which is appropriate for this quantitative study. The report provided evidence that the data collection methods yielded data that were reliable, valid, and responsive.</p>

	data that were reliable, valid and responsive?	
Procedures	<ul style="list-style-type: none"> • If there was an intervention, was it adequately described, and was it rigorously developed and implemented? Did most participants allocated to the intervention group actually receive it? Was there evidence of intervention fidelity? • Were data collected in a manner that minimized bias? Were the staff who collected data appropriately trained? 	<p>The intervention was described as application of intranasal mupirocin and washing with chlorhexidine soap or application of intranasal povidone-iodine and washing with povidone-iodine soap, each treatment lasting for five days. The intranasal mupirocin/chlorhexidine wash group was administered to uncomplicated cases and the povidone-iodine intranasal/soap group was administered to the more complicated cases. The complicated and uncomplicated criteria are listed in a figure in the study. The intervention was more than adequately described and was rigorously developed and implemented. All patients in the intervention group actually received the intervention. Those that did not were excluded from the study. Data were collected in a manner that minimized bias, in a retrospective manner.</p>

Data Analysis	<ul style="list-style-type: none"> • Were analyses undertaken to address each research question or test each hypothesis? • Were appropriate statistical methods used, given the level of measurement of the variables, number of groups being compared, and assumptions of the texts? • Was a powerful analytic method used? (e.g., did the analysis help to control for confounding variables)? • Were type I and Type II errors avoided or minimized? • In intervention studies, was an intention-to-treat analysis performed? • Were problems of missing values evaluated and adequately addressed? 	<p>Analyses were undertaken to address each research question. The appropriate statistical methods were used given the level of measurement of the variables, number of groups being compared, and assumptions of the texts. A powerful analytic method was used described as the Epi Info 7. Normally distributed continuous variables were compared using a 2-sample t test, and categorical data were compared using a X2 test with a Yates correction. Type I and type II errors were avoided. There was no mention of an intention-to-treat analysis being performed. Problems of missing values were evaluated and adequately addressed specifically in this study.</p>
Findings	<ul style="list-style-type: none"> • Was information about statistical significance presented? Was information about effect size and precision of estimates (confidence 	<p>Information about statistical significance was presented. Information about confidence intervals was presented. The findings were adequately summarized with</p>

	<p>intervals) presented?</p> <ul style="list-style-type: none"> • Were the findings adequately summarized, with good use of tables and figures? • Were findings reported in a manner that facilitates a meta-analysis, and with sufficient information needed for EBP? 	<p>appropriate use of tables. The findings were reported in a manner that facilitates a meta-analysis and with sufficient information needed for EBP.</p>
<p>Discussion Interpretation of the Findings</p>	<ul style="list-style-type: none"> • Were all major findings interpreted and discussed within the context of prior research and/or the study's conceptual framework? • Were casual inferences, if any, justified? • Was the issue of clinical significance discussed? • Were interpretations well-founded and consistent with the study's limitations? • Did the report address the issue of the generalizability of the findings? 	<p>All major findings were interpreted and discussed within the context of prior research. The issue of clinical significance was certainly discussed. The interpretations were well-founded and consistent with the study's limitations. The report does address the issue of the generalizability of the findings, stating there was a low rate of successful treatment. This is believed to have been due to the use of povidone-iodine being restricted to only the complicated cases. There were also more patients included in this study with less than ideal prognostic factors such as those with chronic wounds, which are</p>

		contraindicated in most other studies.
Implications/Recommendations	<ul style="list-style-type: none"> Did the researchers discuss the implications of the study for clinical practice or further research—and were those implications reasonable and complete? 	The researchers discussed the implications of the study for clinical practice and they recommend that systematic topical decolonization in MRSA carrier be undertaken in accordance with recent proposals made by other researchers.
General Issues Presentation	<ul style="list-style-type: none"> Was the report well-written, organized, and sufficiently detailed for critical analysis? In intervention studies, was a CONSORT flowchart provided to show the flow of participants in the study? Was the report written in a manner that makes the findings accessible to practicing nurses? 	The report was definitely well-written, organized, and sufficiently detailed for critical analysis. A CONSORT flowchart was provided to show the flow of participants in the study. The report was definitely written in a manner that makes the findings accessible to practicing nurses.
Researcher Credibility	<ul style="list-style-type: none"> Do the researchers' clinical, substantive, or methodologic qualifications and experience enhance confidence in the 	While there is no mention of the researcher's clinical, substantive, or methodologic qualifications and experience, one of the authors, Sai, worked for the Hospital

	findings and their interpretation?	Epidemiology and Infection Control Unit, so it would seem he has clinical qualifications.
Summary Assessment	<ul style="list-style-type: none"> • Despite any limitations, do the study findings appear to be valid—do you have confidence in the truth value of the results? • Does the study contribute any meaningful evidence that can be used in nursing practice or that is useful to the nursing discipline? 	<p>Despite any limitations, the study findings do appear to be valid, although this study could have benefitted from a larger sample size. This study does contribute to meaningful evidence that can be used in nursing practice and will be useful to the nursing discipline as the authors further demonstrate the effectiveness of decolonization on the infection rates.</p>

Appendix A-5

Peng, H. M., Wang, L. C., Zhai, J. L., Weng, X. S., Feng, B., & Wang, W. (2018). Effectiveness of preoperative decolonization with nasal povidone iodine in chinese patients undergoing elective orthopedic surgery: A prospective cross-sectional study. *Brazilian Journal of Medical and Biological Research*, 51(2), 1-6.

Title	<ul style="list-style-type: none"> Is the title a good one, succinctly suggesting key variables and the study population? 	The title was very informative, suggesting key variables and identifying the study population as Chinese patients.
Abstract	<ul style="list-style-type: none"> Did the abstract clearly and concisely summarize the main features of the report (problem, methods, results, conclusions)? 	The abstract concisely summarized the main features of the report including the problem, methods, results, and conclusions.
Introduction Statement of the Problem	<ul style="list-style-type: none"> Was the problem stated unambiguously, and was it easy to identify? Is the problem statement build a persuasive argument for the new study? Was there a good match between the research problem and the methods used –that is, was a quantitative approach appropriate? 	The problem was stated unambiguously and it was easy to identify. The problem statement does build a persuasive argument for the new study. The authors cited the fact that there was data lacking regarding the prevalence and distribution of MSSA and MRSA in patients undergoing orthopedic surgery in China. A quantitative study was appropriate for the stated research problem.

Hypotheses or Research Questions	<ul style="list-style-type: none"> • Were research questions and/or hypotheses explicitly stated? If not, was their absence justified? • Were questions and hypotheses appropriately worded, with clear specification of key variables and the study population? • Were the questions/hypotheses consistent with existing knowledge? 	<p>The research question was explicitly stated. The question was appropriately worded with clear specification of key variables and the study population. The question was consistent with existing knowledge.</p>
Literature Review	<ul style="list-style-type: none"> • Was the literature review up-to-date and based mainly on primary sources? • Did the review provide a state-of-the-art synthesis of evidence on the problem? • Did the literature review provide a strong basis for the new study? 	<p>The literature was fairly up to date, dating as far back as 2002. There was a state-of-the art synthesis of evidence on the problem provided. The literature review provided a strong basis for the new study citing a lack of similar studies in China.</p>
Conceptual/Theoretical Framework	<ul style="list-style-type: none"> • Were key concepts adequately defined conceptually? • Was a conceptual/theoretical framework articulated—and, if so, was it appropriate? If not, is the absence of a framework justified? • Were the questions/hypotheses consistent with the framework? 	<p>Key concepts were adequately defined conceptually. There was no conceptual/theoretical framework articulated. The absence of the framework was not justified.</p>

Method Protection of Human Rights	<ul style="list-style-type: none"> • Were appropriate procedures used to safeguard the rights of study participants? • Was the study externally reviewed by an IRB/ethics review board? • Was the study designed to minimize risks and maximize benefits to participants? 	Appropriate procedures were used to safeguard the right of study participants and informed consent was obtained from the patients before they were swabbed. The study was approved by the Institutional Ethics Committee of Peking Union Medical College Hospital. The study was designed to minimize risks and maximize benefits to study participants.
Research Design	<ul style="list-style-type: none"> • Was the most rigorous design used, given the study purpose? • Were appropriate comparisons made to enhance interpretability of the findings? • Was the number of data collection points appropriate? • Did the design minimize biases and threats to the internal, construct, and external validity of the study (e.g., was blinding used, was attrition minimized)? 	The most rigorous design was used given the purpose of the study. The appropriate comparisons were made to enhance interpretability of the findings. The number of data collection points were appropriate. Blinding was not used.
Population and Sample	<ul style="list-style-type: none"> • Was the population identified? Was the sample described in sufficient detail? 	The study population was identified and described in

	<ul style="list-style-type: none"> • Was the best possible sampling design used to enhance the sample's representativeness? Were sampling biases minimized? • Was the sample size based on a power analysis? 	<p>sufficient detail. The sampling design used was a prospective cross-sectional study. Patients were selected consecutively from those who were undergoing elective orthopedic surgery between August 2015 and February 2016. The minimum sample size was calculated to be 457 patients in order to detect the prevalence with a 2% precision.</p>
Data Collection and Measurement	<ul style="list-style-type: none"> • Were the operational and conceptual definitions congruent? • Were key variables measured using an appropriate method (e.g., interviews, observations, and so on)? • Were specific instruments adequately described and were they good choices, given the study population and the variables being studied? • Did the report provide evidence that the data collection methods yielded data that were reliable, valid and responsive? 	<p>Operational and conceptual definitions were congruent. Key variables were measured appropriately using data collection as well as a questionnaire completed by each patient in order to evaluate each patients' characteristics. Specific instruments were adequately described and were appropriate choices given the study population.</p>

Procedures	<ul style="list-style-type: none"> • If there was an intervention, was it adequately described, and was it rigorously developed and implemented? Did most participants allocated to the intervention group actually receive it? Was there evidence of intervention fidelity? • Were data collected in a manner that minimized bias? Were the staff who collected data appropriately trained? 	<p>There was an intervention and it was adequately identified and described as well as rigorously developed and implemented. There were 33 patients excluded from the study who did not receive any intervention. There was evidence of intervention fidelity. Data were collected in a manner that minimized bias and the staff who collected the data were appropriately trained.</p>
Data Analysis	<ul style="list-style-type: none"> • Were analyses undertaken to address each research question or test each hypothesis? • Were appropriate statistical methods used, given the level of measurement of the variables, number of groups being compared, and assumptions of the tests? • Was a powerful analytic method used? (e.g., did the analysis help to control for confounding variables)? • Were type I and Type II errors avoided or 	<p>Analyses were undertaken to address the research question. All statistical analyses were performed using the software SPSS. The statistical analyses comparing the pre-operative results were performed using the McNemar test. Type I and type II errors were minimized. There was no mention of an intention-to-treat analysis being performed. Problems of missing values were evaluated and adequately</p>

	<ul style="list-style-type: none"> minimized? In intervention studies, was an intention-to-treat analysis performed? Were problems of missing values evaluated and adequately addressed? 	addressed.
Findings	<ul style="list-style-type: none"> Was information about statistical significance presented? Was information about effect size and precision of estimates (confidence intervals) presented? Were the findings adequately summarized, with good use of tables and figures? Were findings reported in a manner that facilitates a meta-analysis, and with sufficient information needed for EBP? 	Information about statistical significance was presented. There was information about effect size and precision of estimates presented. The findings were adequately summarized but the tables and figures could have used further elaboration. The findings were reported in a manner that facilitated a meta-analysis with sufficient information needed for EBP.
Discussion Interpretation of the Findings	<ul style="list-style-type: none"> Were all major findings interpreted and discussed within the context of prior research and/or the study's conceptual framework? Were casual inferences, if any, justified? Was the issue of clinical significance discussed? Were interpretations 	All major findings were interpreted and discussed within the context of prior research. Casual inferences were justified. The clinical significance was discussed. The interpretations were well-founded and consistent with the study's limitations. The report did

	<p>well-founded and consistent with the study's limitations?</p> <ul style="list-style-type: none"> • Did the report address the issue of the generalizability of the findings? 	<p>address the generalizability of the findings.</p>
Implications/Recommendations	<ul style="list-style-type: none"> • Did the researchers discuss the implications of the study for clinical practice or further research—and were those implications reasonable and complete? 	<p>The researchers did discuss the implications of the study for clinical practice and further research. They suggested that the nasal povidone-iodine swabs should be used to eradicate nasal colonization of MRSA/MSSA and they also suggested that research with a larger cohort of orthopedic surgery patients be performed to determine its efficacy in eradicating MRSA/MSSA colonization.</p>
General Issues Presentation	<ul style="list-style-type: none"> • Was the report well-written, organized, and sufficiently detailed for critical analysis? • In intervention studies, was a CONSORT flowchart provided to show the flow of participants in the study? • Was the report written in a manner that makes the 	<p>The report was very well written, organized, and sufficiently detailed for critical analysis. A CONSORT flowchart was provided to show the flow of participants in the study. The report was written in a manner that made the findings accessible to</p>

	findings accessible to practicing nurses?	practicing nurses.
Researcher Credibility	<ul style="list-style-type: none"> Do the researchers' clinical, substantive, or methodologic qualifications and experience enhance confidence in the findings and their interpretation? 	The credentials of the researchers were not identified, but the study was performed by the department of orthopedic surgery. It can be expected that members would have the experience necessary to enhance confidence in the findings and their interpretations.
Summary Assessment	<ul style="list-style-type: none"> Despite any limitations, do the study findings appear to be valid—do you have confidence in the truth value of the results? Does the study contribute any meaningful evidence that can be used in nursing practice or that is useful to the nursing discipline? 	While the sample size was small and the study was conducted only at one community hospital, the study findings do appear to be valid. Readers can have confidence in the truth value of the results. The study does contain meaningful evidence that can be used in nursing practice.

Appendix A-6

Urias, D. S., Varghese, M., Simunich, T., Morrissey, S., & Dumire, R. (2018).
Preoperative decolonization to reduce infections in urgent lower extremity repairs.
European Journal of Trauma and Emergency Surgery, 44, 787-793.

Title	<ul style="list-style-type: none"> Is the title a good one, succinctly suggesting key variables and the study population? 	The title is good, succinctly suggest key variables as well as the study population.
Abstract	<ul style="list-style-type: none"> Did the abstract clearly and concisely summarize the main features of the report (problem, methods, results, conclusions)? 	The abstract very clearly and concisely summarized the main features of the report including the purpose, methods, results, and conclusions.
Introduction Statement of the Problem	<ul style="list-style-type: none"> Was the problem stated unambiguously, and was it easy to identify? Is the problem statement build a persuasive argument for the new study? Was there a good match between the research problem and the methods used –that is, was a quantitative approach appropriate? 	The problem was very easy to identify and was stated unambiguously as measuring the effectiveness in reducing SSIs in patients undergoing repair of lower extremity fractures. The problem statement and the provided research data built a persuasive argument for the new study. There was a good match between the research problem and the methods used, a quantitative approach was appropriate for this study given it was based on data collection and statistics.

Hypotheses or Research Questions	<ul style="list-style-type: none"> • Were research questions and/or hypotheses explicitly stated? If not, was their absence justified? • Were questions and hypotheses appropriately worded, with clear specification of key variables and the study population? • Were the questions/hypotheses consistent with existing knowledge? 	<p>The research question was explicitly stated. IT was appropriately worded with clear specification of key variables and the study population. The question was certainly consistent with existing knowledge with references made to recent research studies.</p>
Literature Review	<ul style="list-style-type: none"> • Was the literature review up-to-date and based mainly on primary sources? • Did the review provide a state-of-the-art synthesis of evidence on the problem? • Did the literature review provide a strong basis for the new study? 	<p>The literature review was up-to-date with most in the past six years, some going back 14 years, and were based on primary resources. The review did provide a state-of-the-art synthesis of evidence on the problem as well as a strong basis for the new study.</p>
Conceptual/Theoretical Framework	<ul style="list-style-type: none"> • Were key concepts adequately defined conceptually? • Was a conceptual/theoretical framework articulated—and, if so, was it appropriate? If not, is the absence of a framework justified? • Were the questions/hypotheses consistent with the 	<p>Key concepts were adequately defined conceptually in this study. There was no conceptual/theoretical framework utilized in this study. The absence of a framework was justified as this quantitative study had a nice flow and was neatly</p>

	framework?	summarized in various tables.
Method Protection of Human Rights	<ul style="list-style-type: none"> • Were appropriate procedures used to safeguard the rights of study participants? • Was the study externally reviewed by an IRB/ethics review board? • Was the study designed to minimize risks and maximize benefits to participants? 	Appropriate procedures were used to safeguard the rights of study participants. There was no mention of a review by an IRB however, there is mention that since this was a retrospective study, informed consent was not obtained. The study was designed to minimize risks and maximize benefits to participants and the study made note that only minimal personal health information was collected and the data was de-identified.
Research Design	<ul style="list-style-type: none"> • Was the most rigorous design used, given the study purpose? • Were appropriate comparisons made to enhance interpretability of the findings? • Was the number of data collection points appropriate? • Did the design minimize biases and threats to the internal, construct, and external validity of the study (e.g., was blinding used, 	The research design used in this study was a retrospective study at a rural community-based hospital. The was the most rigorous design given the purpose of the study. Appropriate comparisons were made to enhance interpretability of the findings. The number of data collection points was appropriate, but could have been expanded to include

	was attrition minimized)?	pre and post-intervention nasal swabs for MSSA and MRSA. There were not performed to save money but would have provided more information. The design did minimize biases and threats to the internal construct and external validity of the study.
Population and Sample	<ul style="list-style-type: none"> • Was the population identified? Was the sample described in sufficient detail? • Was the best possible sampling design used to enhance the sample's representativeness? Were sampling biases minimized? • Was the sample size based on a power analysis? 	The population was very clearly identified and the sample described in sufficient detail and was summarized in provided tables. The best possible sampling design was used to enhance the sample's representativeness and sampling biases were minimized. There was no mention of the sample size being based on a power analysis.
Data Collection and Measurement	<ul style="list-style-type: none"> • Were the operational and conceptual definitions congruent? • Were key variables measured using an appropriate method (e.g., interviews, observations, and so on)? • Were specific instruments 	The operational and conceptual definitions were congruent in this study. The key variables were measured appropriately using statistical analysis for the quantitative study. Specific instruments were

	<p>adequately described and were they good choices, given the study population and the variables being studied?</p> <ul style="list-style-type: none"> • Did the report provide evidence that the data collection methods yielded data that were reliable, valid and responsive? 	<p>adequately described and were good choices given this study population. The instruments used were descriptive demographic statistics, univariate analyses, logistic regression, chi-squared test, Fisher's exact test, and the Mann-Whitney <i>U</i> test. Categorical variables were analyzed using Fisher's exact test. The report provides some evidence that the data collection methods yielded data that was reliable, valid, and responsive.</p>
Procedures	<ul style="list-style-type: none"> • If there was an intervention, was it adequately described, and was it rigorously developed and implemented? Did most participants allocated to the intervention group actually receive it? Was there evidence of intervention fidelity? • Were data collected in a manner that minimized bias? Were the staff who collected data appropriately 	<p>The intervention in this study was a decolonization protocol of bathing patient with 2% CHG washcloths or Dynahex 4% CHG solution in the pre-intervention group and the bathing protocol as well as a povidone-iodine nasal swab 1 hour before surgical incision in the intervention group. It was adequately described, rigorously developed and</p>

	trained?	implemented. The participants who were allocated to the intervention group did receive the intervention. There was no evidence of intervention fidelity. Data was collected in a manner that minimized bias and the staff was appropriately trained on administered the intervention protocol.
Data Analysis	<ul style="list-style-type: none"> • Were analyses undertaken to address each research question or test each hypothesis? • Were appropriate statistical methods used, given the level of measurement of the variables, number of groups being compared, and assumptions of the texts? • Was a powerful analytic method used? (e.g., did the analysis help to control for confounding variables)? • Were type I and Type II errors avoided or minimized? • In intervention studies, was an intention-to-treat 	<p>There was data analysis undertaken to address the research question. There was mention of statistical methods used and they were appropriate to measure the variables, number of groups being compared, and assumptions of the text. There was a powerful analytic method used: logistic regression; univariate analysis; chi-squares test; Fisher's exact test; and the Mann-Whitney <i>U</i> test. Type I and type II errors were minimized. There was no mention of an intention-to-treat</p>

	<p>analysis performed?</p> <ul style="list-style-type: none"> • Were problems of missing values evaluated and adequately addressed? 	<p>analysis being performed. There were no noted missing values.</p>
Findings	<ul style="list-style-type: none"> • Was information about statistical significance presented? Was information about effect size and precision of estimates (confidence intervals) presented? • Were the findings adequately summarized, with good use of tables and figures? • Were findings reported in a manner that facilitates a meta-analysis, and with sufficient information needed for EBP? 	<p>Information about statistical significance was presented. Information about effect size and confidence intervals were presented. The findings were adequately summarized with good use of tables and figures. The findings were reported in a manner that facilitates a meta-analysis with sufficient information needed for EBP.</p>
Discussion Interpretation of the Findings	<ul style="list-style-type: none"> • Were all major findings interpreted and discussed within the context of prior research and/or the study's conceptual framework? • Were casual inferences, if any, justified? • Was the issue of clinical significance discussed? • Were interpretations well-founded and consistent with the study's limitations? • Did the report 	<p>All major findings were interpreted and discussed within the context of prior research. Casual inferences were justified. Clinical significance was discussed. Interpretations were well-founded and consistent with the study's limitations. The report did address the issue of generalizability of the findings noting that this was at a</p>

	address the issue of the generalizability of the findings?	small hospital, with mostly Caucasian patients, with a small scope of surgeries.
Implications/Recommendations	<ul style="list-style-type: none"> Did the researchers discuss the implications of the study for clinical practice or further research—and were those implications reasonable and complete? 	The researchers did discuss the implications of the study for clinical practice stating that the use of intranasal povidone-iodine is effective as well as cost-effective and further research should be performed on different surgical procedures.
General Issues Presentation	<ul style="list-style-type: none"> Was the report well-written, organized, and sufficiently detailed for critical analysis? In intervention studies, was a CONSORT flowchart provided to show the flow of participants in the study? Was the report written in a manner that makes the findings accessible to practicing nurses? 	The report was well written, organized, and sufficiently detailed for critical analysis. There was no CONSORT flowchart provided to show the flow of participants in the study. The report was written in a manner that makes the findings accessible to practicing nurses.
Researcher Credibility	<ul style="list-style-type: none"> Do the researchers' clinical, substantive, or methodologic qualifications and experience enhance confidence in the findings and their interpretation? 	The researchers' clinical qualifications and experience enhanced confidence in the findings and their interpretation.
Summary Assessment	<ul style="list-style-type: none"> Despite any limitations, do the 	While this was a small study at only

	<p>study findings appear to be valid—do you have confidence in the truth value of the results?</p> <ul style="list-style-type: none">• Does the study contribute any meaningful evidence that can be used in nursing practice or that is useful to the nursing discipline?	<p>one, rural institution, there is confidence in the truth value of the results. This study definitely contributed meaningful evidence that can be used in nursing practice and/or the nursing discipline, proving that nasal povidone-iodine can be efficacious and cost-effective in reducing post-operative SSI's.</p>
--	--	--

Appendix B-1

Phillips, M., Rosenberg, A., Shopsis, B., Cuff, G., Skeete, F., Foti, A., Kraemer, K., Inglima, K., Press, B., & Bosco, J. (2014). Preventing surgical site infections: A randomized, open-label trial of nasal mupirocin ointment and nasal povidone iodine solution. *Infection Control Hosp Epidemiol*, 35(7), 826-832

Purpose	Methods	Results
To test the hypothesis that a one-time application of nasal povidone iodine just prior to surgery would be as effective as twice daily applications of nasal mupirocin during the five days before surgery in preventing SSIs and provide a more convenient option for patients at lower cost.	This was an investigator initiated, prospective, open-label, randomized trial.	<p>-Of those subjects in the intent-to-treat analysis, a deep SSI caused by <i>S. Aureus</i> was noted in 5 of 855 surgeries in the mupirocin group and 1 of 842 surgeries in the povidone iodine group ($p=0.2$).</p> <p>-A deep SSI caused by any pathogen developed in 14 surgeries in the mupirocin group and 6 surgeries in the povidone iodine group ($p=0.1$).</p> <p>-In the per protocol analysis, <i>S. aureus</i> deep SSIs developed in 5 of 763 surgeries in the mupirocin group and 0 of 776 surgeries in the povidone iodine group ($p=0.03$).</p> <p>-The findings suggested that pre-operative nasal povidone iodine with topical chlorhexidine is similar to pre-operative nasal mupirocin with topical chlorhexidine in preventing <i>S. aureus</i> deep SSI after arthroplasty and spine fusion surgery.</p> <p>-The study also determined that</p>

		<p>application of nasal povidone iodine by the care team just prior to surgery may ensure greater compliance.</p> <p>-The study identified that <i>S. aureus</i> colonization pre-operatively was a significant risk factor for subsequent <i>S. aureus</i> SSI.</p> <p>-As povidone iodine costs less than a mupirocin course, the authors noted that povidone iodine provided more value as defined as quality of outcomes divided by cost.</p>
--	--	---

Appendix B-2

Anderson, M. J., David, M. L., Scholz, M., Bull, S. J., Morse, D., Hulse-Stevens, M., & Peterson, M. L. (2015). Efficacy of skin and nasal povidone-iodine preparation against mupirocin-resistant methicillin-resistant staphylococcus aureus and *S. aureus* within the anterior nares. *Antimicrobial Agents and Chemotherapy*, 59(5), 2765-2773.

Purpose	Methods	Results
There were three aims to this study: to demonstrate that povidone-iodine nasal prep is effective at reducing and preventing MRSA infection in an ex vivo model; to show efficacy against Mup MRSA in ex vivo human skin MRSA infection model and to demonstrate reduction in the anterior nares <i>S. aureus</i> bioburden of human subjects.	This was a quantitative study with data collected on the efficacy of povidone-iodine or mupirocin intranasally in the treatment of MRSA on ex vivo porcine vaginal mucosa, ex vivo human skin, and the anterior nares of human subjects.	-The antimicrobial effect of 3M nasal povidone-iodine compared to bactroban nasal was tested on 10 Mup MRSA isolates, both high-level resistance and low-level resistance, using ex vivo porcine vaginal mucosa. One hour following application, the povidone-iodine treated explants had significantly less low-level resistance (n=4) MRSA bacteria than untreated controls and bactroban nasal treated explants ($p < 0.05$) (at 1 hour, 1.63 +/- 0.44 versus 5.30 +/- 0.30 and 5.71 +/- 0.57. There was some regrowth noted at 6 hours but povidone-iodine explants were associated with significantly lower bacterial densities, 2.56 +/- 1.60 and 3.62 +/- 0.50 than untreated controls, 6.60 +/- 0.75 and 7.76 +/- 0.22 or bactroban treated

		<p>explants, 5.08 +/- 0.39 and 5.99 +/- CFU explant).</p> <p>-Similarly, at all three time frames examined, high-level resistance MRSA (n=6) infected explants treated with 3M povidone-iodine had significantly lower bacterial densities than untreated on bactroban nasal treated explants ($p<0.05$) (at 1 hour, 1.55 +/- 0.29 versus 5.68 +/- 0.29 or 6.03 +/- 0.32, at 6 hours, 2.98 +/- 0.23 versus 7.04 +/- 0.26 or 5.99 +/- 0.43, and at 24 hours, 3.24 +/- 0.36 versus 7.66 +/- 0.19 or 6.88 +/- 0.24).</p> <p>-3M nasal povidone-iodine was tested on reducing normal flora of human anterior nares (n=70). The mean baseline level of <i>S. aureus</i> in this study was log₁₀ 4.77 +/- 0.62 CFU. The nares were sampled at 1, 6, and 12 hours following application of 3M povidone-iodine (n=13-18) versus saline control (n=7-9). At all time frames, <i>S. aureus</i> was reduced from baseline in the 3M povidone-iodine subjects ($p<0.05$) (2.3 +/- 1.68 versus 0.86 +/- 0.73 at 1 hour, 2.79 +/- 1.52 versus 0.76 +/- 0.58 at 6 hours, and 2.37 +/- 1.77 versus 0.6 +/- 0.9 at</p>
--	--	--

		<p>12 hours).</p> <p>-The final part of the study evaluated whether treatment with 3M povidone-iodine, betadine ophthalmic, or mupirocin could prevent PVM explants from becoming infected with MRSA. Treatment with povidone-iodine resulted in a reduction in the ability of MRSA to infect explants far superior to the Betadine ophthalmic formulation (log₁₀ 0.00 +/- 0.00 versus 2.34 +/- 0.12 CFU. Explants treated with 2% mupirocin had CFU equivalent to the controls (log₁₀ 4.53 +/- 0.05 CFU/explant (n=4, p=<0.05).</p> <p>-The authors concluded that 3M povidone-iodine showed a significant reduction in <i>S. aureus</i> from the anterior nares of human subjects.</p> <p>-The study authors suggested that due to medication compliance and evolving mupirocin resistance, 3M povidone-iodine should be considered for the reduction of <i>S. aureus</i> SSIs. The application of 3M povidone-iodine is health care provider observed thus improving the compliance rate.</p>
--	--	---

Appendix B-3

Bebko, S. P., Green, D. M., & Awad, S. S. (2015). Effect of a preoperative decontamination protocol on surgical site infections in patients undergoing elective orthopedic surgery with hardware implantation. *JAMA Surgery*, 150(5), 390-395.

Purpose	Methods	Results
The purpose of this study was to examine the effect of a decontamination protocol on SSIs in patients undergoing elective orthopedic surgery with hardware implantation. The decontamination protocol consisted of the application of both chlorhexidine washcloths and oral rinse the night before and the morning of surgery as well as intranasal povidone-iodine once the morning of surgery.	The method for this study was a prospective clinical study for patients undergoing elective orthopedic surgery with hardware implants.	<ul style="list-style-type: none"> -There were 344 patients in the control group and 365 patients in the intervention group. -There were 13 patients in the control group who developed an SSI (3.8%)($p=.02$). -There were 4 patients in the intervention group who developed an SSI (1.1%)($p=.02$). -A significant reduction in the number of SSIs (69.2%) was noted in this study in the intervention group. -Decontamination was an independent protective factor against the development of an SSI (CI 0.08-0.77) ($p=.02$). -The results of the study showed a 100% adherence rate to their 2-day protocol. -The authors found that there was a greater cost savings using nasal povidone-iodine and chlorhexidine washcloths (\$35/patient) versus \$54.72 per patient to use mupirocin and chlorhexidine

		washcloths.
--	--	-------------

Appendix B-4

Sai, N., Laurent, C., Strale, H., Denis, O., & Byl, B. (2015). Efficacy of the decolonization of methicillin-resistant staphylococcus aureus carriers in clinical practice. *Antimicrobial Resistance and Infection Control*, 4(56), 1-8.

Purpose	Methods	Results
The purpose of this study was to evaluate the effectiveness of two decolonization protocols for newly diagnosed MRSA colonization in hospitalized patients and to assess the impact of decolonization on the rate of MRSA infection.	<p>-This was a retrospective cohort study to test two decolonization protocols on the rate of MRSA infection in patients newly diagnosed as colonized with MRSA.</p> <p>-Study participants received one of two decolonization protocols: the application of intranasal mupirocin 2% and washing with chlorhexidine soap or application of intranasal povidone-iodine and washing with povidone-iodine soap, with each treatment lasting for 5 days.</p>	<p>- There were 268 patients who were eligible for the decontamination protocol in this study.</p> <p>-39% of patients were successfully decolonized whereas 164 were not.</p> <p>-Those treated with povidone-iodine intranasally had a 18% success rate in the first attempt and a 16% success rate in the second attempt at decolonization.</p> <p>-Those treated with mupirocin intranasally had a 51% success rate in the first treatment attempt and a 29% success rate in the second attempt.</p> <p>-The total success rate for mupirocin was 56% and the total success rate for povidone-iodine was 23%.</p> <p>-The study showed that only one-third of patients were successfully decolonized after the first treatment attempt, with 39% decolonized following the second attempt, with no further successes beyond the first two</p>

		<p>attempts in those with previous failures.</p> <p>-Of those who were successfully decolonized (n=104), none had a subsequent MRSA infection in a median time frame of 43 days. Of those unsuccessfully decolonized (n=164), 8 had a subsequent MRSA infection in the median time frame of 43 days.</p> <p>-This study had a low rate of successful treatment compared to other studies, which the authors attributed to differences in prevalence of failure risk factors and a small sample size. Despite this low success rate, the authors concluded that the effectiveness of decolonization on the infection rate justified the continuation of their strategy.</p>
--	--	--

Appendix B-5

Peng, H. M., Wang, L. C., Zhai, J. L., Weng, X. S., Feng, B., & Wang, W. (2018). Effectiveness of preoperative decolonization with nasal povidone iodine in Chinese patients undergoing elective orthopedic surgery: A prospective cross-sectional study. *Brazilian Journal of Medical and Biological Research*, 51(2), 1-6.

Purpose	Methods	Results
The purpose of this study was to assess the prevalence of MSSA/MRSA in the patients admitted to the study institution and secondarily to determine whether the current treatment protocols result in the successful decolonization of MSSA/MRSA.	<p>-This was a prospective, cross-sectional study.</p> <p>-Patients were screened for nasal MRSA/MSSA colonization within 24 hours of admission.</p> <p>Those positive for MSSA and/or MRSA underwent the decolonization procedure which was povidone-iodine swabs in both nostrils twice a day for five days prior to surgery as well as chlorhexidine gluconate baths for the five days prior to surgery.</p>	<p>-Out of the 545 patients included in this study, there was a total of 72 patients who were positive for MSSA or MRSA who received the decolonization treatment, which consisted of swabbing the nares with 5% povidone-iodine nasal swabs in both nostrils twice a day for 5 days prior to surgery.</p> <p>-The decolonization was 100% successful in eradicating MRSA ($p < 0.000$) and there was a 94% reduction in MSSA colonization ($p < 0.001$).</p> <p>-The compliance rate for the decolonization protocol was noted to be 98.4%.</p> <p>-The study concluded that the nasal treatment with povidone-iodine was as effective as nasal mupirocin.</p>

Appendix B-6

Urias, D. S., Varghese, M., Simunich, T., Morrissey, S., & Dumire, R. (2018).
Preoperative decolonization to reduce infections in urgent lower extremity repairs.
European Journal of Trauma and Emergency Surgery, 44, 787-793.

Purpose	Methods	Results
The purpose of this study was to measure the effectiveness of a CHG bath preoperatively as well as nasal painting using povidone-iodine skin and nasal antiseptic in reducing SSIs in patients undergoing orthopedic operations conducted at a rural hospital.	<p>-This was a retrospective review of trauma patients undergoing orthopedic operations at a rural medical center from 2012-2016 with the intervention period being from 2014-2016.</p> <p>-The CHG baths were routinely performed from October 2012-September 2016. From October 2014-September 2016, CHG baths and nasal povidone-iodine were routinely performed, this was the intervention group.</p>	<p>-This study was performed from October 1, 2012 to September 30, 2016 with a total of 1,746 trauma patients undergoing 1892 orthopedic surgeries to repair fractures of the lower extremities using hardware.</p> <p>-There were 862 patients in the pre-intervention group and 884 patients in the intervention group.</p> <p>-There were a total of 10 SSI's in the pre-intervention group (n=930) and 2 SSI's in the intervention group (n=962) (p=0.020). Of these SSIs, 2 were MRSA in the pre-intervention and intervention groups and 8 were MRSA in the pre-intervention group, zero being MRSA in the intervention group.</p> <p>-There was a significant decrease in the SSI rate from 1.1% in the pre-intervention group to 0.2% in the intervention group (p=0.020).</p> <p>-The pre-intervention annual infection rate for</p>

		<p>the first year of the study was 1.5% (n=476), the second year was 0.7% (n=454). The intervention annual infection rate for the first year of the study was 0.2% (n=484) and 0.2% for the second year (n=478).</p> <p>-The study results indicated that the results of this study have statistical significance, the protocol is cost effective, easy to use, and povidone-iodine is widely available. The researchers encourage multi-center studies and are employing this protocol to other surgical specialties at their institution.</p>
--	--	---

Appendix C

Cross Study Analysis

Author	Phillips et al. (2014).
Key Findings	<p>-The use of nasal povidone-iodine pre-operatively with topical chlorhexidine is similar in efficacy to using nasal mupirocin pre-operatively with topical chlorhexidine in preventing <i>S. aureus</i> deep SSI after arthroplasty and spine fusion surgery.</p> <p>-There was a statistically significant reduction in <i>S. aureus</i> deep SSI in the per protocol group, 5/763 cases in the mupirocin group and 0/776 cases in the povidone-iodine group ($p=0.03$).</p> <p>-Application of nasal povidone-iodine by the healthcare worker just prior to surgery may ensure greater compliance.</p> <p>-Given that this study found povidone-iodine and mupirocin to be equally effective, povidone iodine is the more cost-effective option.</p>
Recommendations	<p>-Target enrollment was not met, but researchers feel study was large enough that a statistical difference was noted.</p> <p>-This study was performed at one institution and could use further studies in different patient populations.</p> <p>-Nasal povidone-iodine has proven to be considered as an alternative to mupirocin to reduce SSIs.</p>
Author	Anderson et al. (2015).
Key Findings	<p>-When applied to the anterior nares of human test subjects, intranasal povidone-iodine rapidly achieved a significant reduction of <i>S. aureus</i> ($p<0.05$).</p> <p>-When the efficacy of povidone-iodine was tested on MRSA-infected ex vivo human skin, CFU were significantly lower than the untreated controls. Zero CFU were recovered from the povidone-iodine treated explants at 12 hours post wash. Intranasal povidone-iodine was</p>

	significantly more effective at reducing MRSA than Betadine Ophthalmic or Bactroban Nasal ($p<0.05$).
Recommendations	<ul style="list-style-type: none"> -The benefits of intranasal povidone-iodine should be considered given issues with medication compliance and evolving mupirocin resistance. -The benefits of intranasal povidone-iodine include: rapid efficacy; broad-spectrum activity against multiple opportunistic pathogens; lack of development of antimicrobial resistance; ease of use; and is directly provided by a health care provider.
Author	Bebko et al. (2015).
Key Findings	<ul style="list-style-type: none"> -This study noted a significant decrease in SSIs after implementing a decontamination protocol in patients undergoing elective orthopedic surgery with hardware. The decontamination protocol consisted of chlorhexidine washcloths, chlorhexidine oral rinse, and intranasal povidone-iodine. In the control group 3.8% of the patients developed an SSI ($n=344$). In the intervention group, 1.1% of patient developed a SSI ($n=365$). -Methicillin resistant staphylococcus aureus decontamination was found to be an independent predictor of not developing an SSI (95% CI, $p=.02$).
Recommendations	<ul style="list-style-type: none"> -This was a study performed on mostly male veterans which may undermine the external validity. -The follow-up period for SSI detection was limited to 30 days which could have affected the sensitivity of SSI identification. -The researchers felt that their data demonstrated a significant decrease in SSI rates after the implementation of their decontamination protocol and recommended further, large-scale randomized controlled clinical trials to be implemented in the future.
Author	Sai et al. (2015).

Key Findings	<p>-In this study, 39% of study participants were successfully decolonized for MRSA, 164 were not. Of those treated with nasal mupirocin/CHG wash the success rate was 51% MRSA reduction in the first attempt and 16% in the second attempt. Of those treated with povidone-iodine and intranasal povidone-iodine, there was an 18% decolonization of MRSA in the first attempt and 23% in the second attempt.</p> <p>-The success rates of mupirocin/CHG was 56% and the success rate for povidone-iodine was 23%.</p> <p>-Of those successfully decolonized (n=104), there were zero subsequent MRSA infections. Of the unsuccessfully decolonized (n=164), there were 8 subsequent MRSA infections.</p>
Recommendations	<p>-Though those participants treated with povidone-iodine had lower success rates at eradicating MRSA than those treated with mupirocin, the povidone-iodine was applied to the complicated cases, whereas mupirocin was applied to the uncomplicated cases.</p> <p>-The low rate of successful treatment may be reflected by differences in prevalence of failure risk factors.</p> <p>-The researchers suggest that an oral decolonization component be added to the decolonization procedures to improve the result of decolonization.</p> <p>-While the rate of decolonization was not high, the effectiveness of decolonization on the infection rate justifies continuation of this strategy.</p>
Author	Peng et al. (2018).
Key Findings	<p>-Pre-decolonization screening showed 64 patients were positive for MSSA (n=72) and 8 patients were positive for MRSA (n=72). After receiving nasal povidone-iodine and a daily CHG bath for five days patients were re-swabbed and 3 were positive for MSSA (0.6%) and zero were positive for MRSA.</p> <p>-There was a 94% reduction in MSSA</p>

	<p>colonization and a 100% reduction in MRSA colonization.</p> <p>-The use of nasal povidone-iodine is more cost effective than intranasal mupirocin.</p>
Recommendations	<p>-This study used a culture swab to test for MRSA/MSSA rather than by PCR which could increase the sensitivity of detection.</p> <p>-The treatment of nasal povidone-iodine proved to be just as effective as intranasal mupirocin and is more cost effective.</p> <p>-The researchers suggest nasal povidone-iodine be evaluated in larger cohorts of orthopedic surgery patients to determine its efficacy in eradicating MRSA/MSSA colonization to reduce SSIs.</p>
Author	Urias et al. (2018).
Key Findings	<p>-Of those patients in the pre-intervention group (n=930), who were treated with CHG bath, 1.1% developed an SSI post-operatively.</p> <p>-In the intervention group (n=962), participants were treated with CHG bath and intranasal povidone-iodine, 0.2% developed an SSI post-operatively.</p>
Recommendations	<p>-The generalizability of the results may be limited due to the retrospective design, the focused trauma population, lack of ethnic diversity, and location.</p> <p>-The results of this study met statistical significance, the intervention is cost effective, easy to use, and the products are widely available.</p> <p>-The researchers encourage multi-center studies to test the effectiveness of the decolonization protocol in different patient populations.</p>